

Calvin University

Calvin Digital Commons

University Faculty Publications

University Faculty Scholarship

12-1-2009

Geometry of control-affine systems

Jeanne N. Clelland

University of Colorado Boulder

Christopher G. Moseley

Calvin University

George R. Wilkens

University of Hawai'i at Mānoa

Follow this and additional works at: https://digitalcommons.calvin.edu/calvin_facultypubs



Part of the [Algebraic Geometry Commons](#)

Recommended Citation

Clelland, Jeanne N.; Moseley, Christopher G.; and Wilkens, George R., "Geometry of control-affine systems" (2009). *University Faculty Publications*. 280.

https://digitalcommons.calvin.edu/calvin_facultypubs/280

This Article is brought to you for free and open access by the University Faculty Scholarship at Calvin Digital Commons. It has been accepted for inclusion in University Faculty Publications by an authorized administrator of Calvin Digital Commons. For more information, please contact dbm9@calvin.edu.

Geometry of Control-Affine Systems^{*}

Jeanne N. CLELLAND[†], Christopher G. MOSELEY[‡] and George R. WILKENS[§]

[†] Department of Mathematics, 395 UCB, University of Colorado, Boulder, CO 80309-0395, USA
E-mail: Jeanne.Clelland@colorado.edu

[‡] Department of Mathematics and Statistics, Calvin College, Grand Rapids, MI 49546, USA
E-mail: cgm3@calvin.edu

[§] Department of Mathematics, University of Hawaii at Manoa, 2565 McCarthy Mall,
Honolulu, HI 96822-2273, USA
E-mail: grw@math.hawaii.edu

Received April 02, 2009, in final form September 28, 2009; Published online October 07, 2009
doi:[10.3842/SIGMA.2009.095](https://doi.org/10.3842/SIGMA.2009.095)

Abstract. Motivated by control-affine systems in optimal control theory, we introduce the notion of a *point-affine distribution* on a manifold \mathcal{X} – i.e., an affine distribution \mathcal{F} together with a distinguished vector field contained in \mathcal{F} . We compute local invariants for point-affine distributions of constant type when $\dim(\mathcal{X}) = n$, $\text{rank}(\mathcal{F}) = n - 1$, and when $\dim(\mathcal{X}) = 3$, $\text{rank}(\mathcal{F}) = 1$. Unlike linear distributions, which are characterized by integer-valued invariants – namely, the rank and growth vector – when $\dim(\mathcal{X}) \leq 4$, we find local invariants depending on arbitrary functions even for rank 1 point-affine distributions on manifolds of dimension 2.

Key words: affine distributions; control theory; exterior differential systems; Cartan’s method of equivalence

2000 Mathematics Subject Classification: 58A30; 53C17; 58A15; 53C10

1 Introduction

In [8] and [9], we introduced the notion of *sub-Finsler geometry* as a natural geometric setting for certain problems in optimal control theory. In local coordinates, a control system may be represented as an underdetermined system of ordinary differential equations of the form

$$\dot{x} = f(x, u), \tag{1.1}$$

where $x \in \mathbb{R}^n$ represents the *state vector* of the system and $u \in \mathbb{R}^s$ represents the *control vector*, i.e., variables which may be specified freely in order to “steer” the system in a desired direction. More generally, x and u may take values in an n -dimensional manifold \mathcal{X} and an s -dimensional manifold \mathcal{U} , respectively. Typically there are constraints on how the system may be “steered” from one state to another, so that $s < n$. The systems of greatest interest are *controllable*, i.e., given any two states x_1, x_2 , there exists a solution curve of (1.1) connecting x_1 to x_2 .

The control systems for which sub-Finsler (or sub-Riemannian) geometry are most relevant are *control-linear* systems. These are systems for which the right-hand side of (1.1) is linear in the control variables u and depends smoothly on the state variables x ; i.e., systems of the form

$$\dot{x} = A(x)u, \tag{1.2}$$

where $A(x)$ is an $n \times s$ matrix whose entries are smooth functions of x . For such a system, *admissible* paths in the state space are those for which the tangent vector to the path at each

^{*}This paper is a contribution to the Special Issue “Élie Cartan and Differential Geometry”. The full collection is available at <http://www.emis.de/journals/SIGMA/Cartan.html>

point $x \in \mathcal{X}$ is contained in the subspace $\mathcal{D}_x \subset T_x\mathcal{X}$ determined by the image of the matrix $A(x)$. If this matrix has constant (maximal) rank s on \mathcal{X} , then \mathcal{D} is a rank s linear distribution¹ on \mathcal{X} . Thus the admissible paths in the state space are precisely the *horizontal curves* of the distribution \mathcal{D} , i.e., curves whose tangent vectors at each point are contained in \mathcal{D} .

A sub-Finsler (resp., sub-Riemannian) metric on the distribution $(\mathcal{X}, \mathcal{D})$ is defined by specifying a Finsler (resp., Riemannian) metric on each of the subspaces \mathcal{D}_x . Such a metric may represent a cost function for the control problem, and geodesics for the metric may be thought of as stationary trajectories for the corresponding *optimal control* problem.

While the class (1.2) of control-linear systems contains many interesting examples, a much broader class of interest is that of *control-affine* systems, i.e., systems of the form

$$\dot{x} = a_0(x) + A(x)u, \quad (1.3)$$

where $A(x)$ is a smoothly-varying $n \times s$ matrix and $a_0(x)$ is a smooth vector field on \mathcal{X} . (The vector field $a_0(x)$ is known as the *drift* vector field.)

Example 1.1. The class of control-affine systems contains as a sub-class the *linear* control systems. These are systems of the form

$$\dot{x} = Ax + Bu, \quad (1.4)$$

where A is a constant $n \times n$ matrix and B is a constant $n \times s$ matrix. Note that despite the terminology, the linear control system (1.4) is not control-linear unless $A = 0$.

The geometric structure corresponding to a control-affine system is an *affine distribution* on the state space \mathcal{X} . More precisely (see [12, 13]):

Definition 1.2. A *rank s affine distribution* \mathcal{F} on an n -dimensional manifold \mathcal{X} is a smoothly-varying family of s -dimensional, affine linear subspaces $\mathcal{F}_x \subset T_x\mathcal{X}$. We will say that \mathcal{F} is *strictly affine* if none of the affine subspaces $\mathcal{F}_x \subset T_x\mathcal{X}$ are linear subspaces. Associated to an affine distribution \mathcal{F} is the *direction distribution*

$$L_{\mathcal{F}} = \{\xi_1 - \xi_2 \mid \xi_1, \xi_2 \in \mathcal{F}\}.$$

Note that $L_{\mathcal{F}}$ is a rank s linear distribution on \mathcal{X} .

In this paper, we will limit our attention to distributions (both linear and affine) which are *bracket-generating* (or *almost bracket-generating* in the affine case) and have *constant type*. In order to make these notions precise, we need the notion of the *growth vector* of a distribution (cf. [21]). Let \mathcal{D} be a linear distribution on a manifold \mathcal{X} , and let \mathcal{D} also denote the sheaf of smooth vector fields on \mathcal{X} which are local sections of \mathcal{D} . The iterated Lie brackets of vector fields in \mathcal{D} generate a flag of subsheaves

$$\mathcal{D} = \mathcal{D}^1 \subset \mathcal{D}^2 \subset \cdots \subset T\mathcal{X},$$

defined by $\mathcal{D}^1 = \mathcal{D}$, and

$$\mathcal{D}^{i+1} = \mathcal{D}^i + [\mathcal{D}, \mathcal{D}^i], \quad i \geq 1.$$

At each point $x \in \mathcal{X}$, this flag of subsheaves gives a flag of subspaces of $T_x\mathcal{X}$:

$$\mathcal{D}_x \subset \mathcal{D}_x^2 \subset \cdots \subset T_x\mathcal{X}.$$

¹We use the adjective “linear” to distinguish the usual notion of a distribution \mathcal{D} , where each fiber \mathcal{D}_x is a linear subspace of $T_x\mathcal{X}$, from that of an affine distribution \mathcal{F} . Unless otherwise specified, the word “distribution” will refer to a linear distribution throughout this paper.

Definition 1.3. Let $\mathcal{D}^\infty = \cup_{i \geq 1} \mathcal{D}^i \subset T\mathcal{X}$.

- The smallest integer $r = r(x)$ such that $\mathcal{D}_x^r = \mathcal{D}_x^\infty$ is called the *step* of the distribution at x .
- The distribution \mathcal{D} is *bracket-generating* if $\mathcal{D}^\infty = T\mathcal{X}$.
- Set $n_i(x) = \dim \mathcal{D}_x^i$. The *growth vector* of \mathcal{D} at x is the integer list $(n_1(x), n_2(x), \dots, n_r(x))$, where $r(x)$ is the step of \mathcal{D} at x .
- The distribution \mathcal{D} has *constant type* if $n_i(x)$ is constant on \mathcal{X} for all i ; i.e., if the growth vector of \mathcal{D} is constant on \mathcal{X} .

Remark 1.4. When working with distributions of constant type, it is customary to consider only bracket-generating distributions. For if \mathcal{D} is not bracket-generating, then \mathcal{D}^∞ is a Frobenius distribution on \mathcal{X} . The integral manifolds of \mathcal{D}^∞ define a local foliation of \mathcal{X} with the property that any horizontal curve of \mathcal{D} is contained in a single leaf of the foliation, and the restriction of \mathcal{D} to each leaf of the foliation is bracket-generating.

We can define similar notions for affine distributions: let \mathcal{F} be an affine distribution on a manifold \mathcal{X} , with direction distribution $L_{\mathcal{F}}$. Let \mathcal{F} also denote the sheaf of smooth vector fields on \mathcal{X} which are local sections of \mathcal{F} . The flag of subsheaves

$$\mathcal{F} = \mathcal{F}^1 \subset \mathcal{F}^2 \subset \dots \subset T\mathcal{X}$$

may be defined in the same manner as for distributions: set $\mathcal{F}^1 = \mathcal{F}$, and

$$\mathcal{F}^{i+1} = \mathcal{F}^i + [\mathcal{F}, \mathcal{F}^i], \quad i \geq 1.$$

At each point $x \in \mathcal{X}$, this gives a flag of affine subspaces of $T_x\mathcal{X}$:

$$\mathcal{F}_x \subset \mathcal{F}_x^2 \subset \dots \subset T_x\mathcal{X}.$$

Each affine distribution \mathcal{F}^i in the flag has an associated direction distribution $L_{\mathcal{F}^i}$; these distributions clearly have the property that

$$L_{\mathcal{F}}^i \subset L_{\mathcal{F}^i}.$$

The step, the growth vector, and the notion of bracket-generating are defined for affine distributions in the same way as for linear distributions. But we will also want to consider affine distributions with a slightly weaker bracket-generating property.

Definition 1.5. An affine distribution \mathcal{F} on an n -dimensional manifold \mathcal{X} is *almost bracket-generating* if $\text{rank}(\mathcal{F}^\infty) = n - 1$, and for each $x \in \mathcal{X}$ and any $\xi(x) \in \mathcal{F}_x$, $\text{span}(\xi(x), (L_{\mathcal{F}^\infty})_x) = T_x\mathcal{X}$.

We also impose an additional condition in the definition of constant type – namely, that \mathcal{F} is strictly affine, and that each element \mathcal{F}^i is either strictly affine or a linear distribution. This condition is reflected in the second condition of the following definition.

Definition 1.6. The affine distribution \mathcal{F} has *constant type* if

- $n_i(x)$ is constant on \mathcal{X} for all i ; i.e., the growth vector of \mathcal{F} is constant on \mathcal{X} , and
- for any section ξ of \mathcal{F} , $\dim(\text{span}(\xi(x), (L_{\mathcal{F}^i})_x))$ is constant on \mathcal{X} for all i .

The well-known theorems of Frobenius, Pfaff, and Engel (cf. Section 2.1) imply that linear distributions of constant type on manifolds of dimension $n \leq 4$ have only integer-valued invariants: specifically, any two linear distributions of constant type on manifolds of the same dimension, with the same rank and growth vector, are locally equivalent via a diffeomorphism of the underlying manifolds. Beginning with $n = 5$ and $s = 2$, local invariants depending on arbitrary functions appear: the famous paper of É. Cartan [7] describes local invariants of rank 2 distributions on 5-manifolds with growth vector $(2, 3, 5)$. More recently, Bryant [5] has described local invariants of rank 3 distributions on 6-manifolds with growth vector $(3, 6)$, and Doubrov and Zelenko [10, 11] have given a fairly comprehensive treatment of maximally nonholonomic distributions of ranks 2 and 3 on manifolds of arbitrary dimension.

The study of affine distributions is more recent and less extensive. Elkin [12, 13] has studied equivalence of affine distributions on manifolds of dimension $n \leq 4$ under diffeomorphisms of the underlying manifolds, resulting in local normal forms for the associated control systems. Local invariants appear in lower dimensions than for linear distributions: the first local invariants arise in the case of rank 1 affine distributions on 3-dimensional manifolds. Local normal forms for generic rank 1 affine distributions on \mathbb{R}^3 have also been obtained by Agrachev [1] and Wilkens [25], and for generic rank 1 affine distributions on \mathbb{R}^n and generic rank 2 affine distributions on \mathbb{R}^4 and \mathbb{R}^5 by Agrachev and Zelenko [3].

Given a control-affine system (1.3), one can canonically associate to it an affine distribution \mathcal{F} on the state space \mathcal{X} : if we let $a_1(x), \dots, a_s(x)$ denote the columns of the matrix $A(x)$, then

$$\mathcal{F}_x = \left\{ a_0(x) + \sum_{i=1}^s u^i a_i(x) \mid u^1, \dots, u^s \in \mathbb{R} \right\}.$$

Elkin defines two systems

$$\dot{x} = a_0(x) + \sum_{i=1}^s u^i a_i(x), \tag{1.5}$$

$$\dot{y} = b_0(y) + \sum_{i=1}^s v^i b_i(y) \tag{1.6}$$

on state spaces \mathcal{X} , \mathcal{Y} , respectively, to be *affine equivalent* if there exists a diffeomorphism $\psi : \mathcal{X} \rightarrow \mathcal{Y}$ such that any absolutely continuous curve $x(t)$ in \mathcal{X} is a solution of (1.5) if and only if the curve $y(t) = \psi(x(t))$ is a solution of (1.6). More specifically, such a diffeomorphism must have the property that

$$\begin{aligned} \psi_*(a_0(x)) &= b_0(\psi(x)) + \sum_{j=1}^s \lambda_0^j(x) b_j(\psi(x)), \\ \psi_*(a_i(x)) &= \sum_{j=1}^s \lambda_i^j(x) b_j(\psi(x)), \quad 1 \leq i \leq s \end{aligned} \tag{1.7}$$

for some functions λ_0^j, λ_i^j on \mathcal{X} . In terms of the associated affine distributions $\mathcal{F}_x, \mathcal{F}_y$, this is equivalent to the statement that $\psi_*(\mathcal{F}_x) = \mathcal{F}_y$.

This notion of equivalence is quite natural, especially in terms of the geometry of affine distributions. But with an eye towards studying *optimal* control problems of the form (1.3), we propose a slightly more restrictive definition of equivalence. According to (1.7), the drift vector field $a_0(x)$ in (1.3) may be replaced by any vector field of the form $a_0(x) + \sum_{j=1}^s \lambda_0^j(x) a_j(x)$, and the resulting control system will be affine equivalent to the original. But in practice, there is often a preferred choice for the drift vector field, corresponding to a zero value for some physical

control inputs. This is particularly true in optimal control, where there is typically a specific control input whose cost function is minimal. In this paper, therefore, we will study the following type of geometric object, corresponding to a control-affine system (1.3) with a fixed drift vector field a_0 :

Definition 1.7. A *point-affine distribution* on a manifold \mathcal{X} is an affine distribution \mathcal{F} on \mathcal{X} , together with a distinguished vector field $a_0 \in \mathcal{F}$.

We will consider the equivalence problem for point-affine distributions and their associated control-affine systems with respect to the following notion of equivalence:

Definition 1.8. The point-affine distributions

$$\mathcal{F}_{\mathcal{X}} = a_0 + \text{span}(a_1, \dots, a_s), \quad \mathcal{F}_{\mathcal{Y}} = b_0 + \text{span}(b_1, \dots, b_s)$$

on the manifolds \mathcal{X}, \mathcal{Y} (corresponding to the control-affine systems (1.5) and (1.6), respectively) will be called *point-affine equivalent* if there exists a diffeomorphism $\psi : \mathcal{X} \rightarrow \mathcal{Y}$ such that

$$\begin{aligned} \psi_*(a_0(x)) &= b_0(\psi(x)), \\ \psi_*(a_i(x)) &= \sum_{j=1}^s \lambda_i^j(x) b_j(\psi(x)), \quad 1 \leq i \leq s \end{aligned}$$

for some functions λ_i^j on \mathcal{X} .

In other words, there must exist a diffeomorphism that preserves both the affine distributions *and* the distinguished vector fields.

The primary goal of this paper is the local classification of point-affine distributions of constant type with respect to point-affine equivalence. We will use Cartan's method of equivalence to compute local invariants for point-affine distributions. The remainder of the paper is organized as follows: in Section 2, we review some well-known results on the geometry of linear distributions of constant type, as well as some of Elkin's results on affine distributions, in order to put our results in context. In Section 3 we compute local invariants for point-affine distributions of constant type when $\dim(\mathcal{X}) = n$, $\text{rank}(\mathcal{F}) = n - 1$, and when $\dim(\mathcal{X}) = 3$, $\text{rank}(\mathcal{F}) = 1$. In Section 4 we discuss examples and potential directions for further study.

2 Review of prior results

2.1 Classical normal form results for linear distributions

The following results are well-known; see, e.g., [4] or [16]. We include them here primarily to provide historical context for our results; some of them will also be used in the proofs in Section 3.

Definition 2.1. A distribution \mathcal{D} on a manifold \mathcal{X} is called *completely integrable* (or, more succinctly, *integrable*), if $\mathcal{D}^\infty = \mathcal{D}$.

Theorem 2.2 (Frobenius). *Let \mathcal{D} be a rank s distribution on an n -dimensional manifold \mathcal{X} . If \mathcal{D} is completely integrable, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, \dots, x^n) such that*

$$\mathcal{D} = \text{span} \left(\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^s} \right),$$

or, equivalently,

$$\mathcal{D}^\perp = \{dx^{s+1}, \dots, dx^n\}.$$

Theorem 2.3 (Pfaff). *Let \mathcal{D} be a rank n distribution on an $(n + 1)$ -dimensional manifold \mathcal{X} , and let θ be a nonvanishing 1-form on \mathcal{X} such that $\mathcal{D} = \{\theta\}^\perp$. Let k be the integer defined by the conditions*

$$\theta \wedge (d\theta)^k \neq 0, \quad \theta \wedge (d\theta)^{k+1} = 0.$$

(k is called the Pfaff rank of θ .) *In a sufficiently small neighborhood of any point $x \in \mathcal{X}$ on which k is constant, there exist local coordinates (x^0, \dots, x^n) such that*

$$\theta = \begin{cases} dx^1 + x^2 dx^3 + \dots + x^{2k} dx^{2k+1} & \text{if } (d\theta)^{k+1} = 0, \\ x^0 dx^1 + x^2 dx^3 + \dots + x^{2k} dx^{2k+1} & \text{if } (d\theta)^{k+1} \neq 0. \end{cases}$$

Theorem 2.4 (Engel). *Let \mathcal{D} be a rank 2 distribution of constant type on a 4-dimensional manifold \mathcal{X} , with growth vector $(2, 3, 4)$. Then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3, x^4) such that*

$$\mathcal{D}^\perp = \{dx^2 - x^3 dx^1, dx^3 - x^4 dx^1\}.$$

2.2 Local classification of rank 1 affine systems on 2- and 3-dimensional manifolds under affine equivalence

Linear distributions of rank 1 have no local invariants, as all nonvanishing vector fields on a manifold are well-known to be locally equivalent. Affine distributions, however, may have local invariants even in rank 1. The following results are due to Elkin [12, 13].

Theorem 2.5 (Elkin). *Let \mathcal{F} be a rank 1 strictly affine distribution of constant type on a 2-dimensional manifold \mathcal{X} .*

1. *If \mathcal{F} is almost bracket-generating, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2) such that*

$$\mathcal{F} = \frac{\partial}{\partial x^1} + \text{span} \left(\frac{\partial}{\partial x^2} \right).$$

2. *If \mathcal{F} is bracket-generating, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2) such that*

$$\mathcal{F} = x^2 \frac{\partial}{\partial x^1} + \text{span} \left(\frac{\partial}{\partial x^2} \right).$$

Note that, for reasons of dimension, any strictly affine distribution of rank 1 on a 2-manifold is either almost bracket-generating or bracket-generating.

Theorem 2.6 (Elkin). *Let \mathcal{F} be a rank 1 strictly affine distribution of constant type on a 3-dimensional manifold \mathcal{X} .*

1. *If \mathcal{F} is almost bracket-generating, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3) such that*

$$\mathcal{F} = \left(\frac{\partial}{\partial x^1} + x^3 \frac{\partial}{\partial x^2} \right) + \text{span} \left(\frac{\partial}{\partial x^3} \right).$$

2. If \mathcal{F} is bracket-generating and $L_{\mathcal{F}^2}$ (which must have rank 2) is Frobenius, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3) such that

$$\mathcal{F} = \left(x^2 \frac{\partial}{\partial x^1} + x^3 \frac{\partial}{\partial x^2} \right) + \text{span} \left(\frac{\partial}{\partial x^3} \right).$$

3. If \mathcal{F} is bracket-generating and $L_{\mathcal{F}^2}$ is not Frobenius, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3) such that

$$\mathcal{F} = \frac{\partial}{\partial x^1} + \text{span} \left(x^3 \frac{\partial}{\partial x^1} + \frac{\partial}{\partial x^2} + H \frac{\partial}{\partial x^3} \right),$$

where H is an arbitrary function on \mathcal{X} satisfying $\frac{\partial H}{\partial x^1} \neq 0$.

Note that if a rank 1 strictly affine distribution \mathcal{F} on a 3-manifold is not bracket-generating or almost bracket-generating, then $\text{span}(\mathcal{F})$ is a rank 2 Frobenius distribution $\mathcal{D}_{\mathcal{F}}$ on \mathcal{X} , and \mathcal{F} may be regarded (at least locally) as living on the 2-dimensional leaves of the foliation which is tangent to $\mathcal{D}_{\mathcal{F}}$.

3 Local point-affine equivalence results for point-affine distributions

3.1 Equivalence problem for rank 1 point-affine distributions on 2-manifolds

For point-affine equivalence, the first local invariants appear in the simplest nontrivial case: rank 1 point-affine distributions on 2-dimensional manifolds.

Theorem 3.1. *Let \mathcal{F} be a rank 1 strictly affine point-affine distribution of constant type on a 2-dimensional manifold \mathcal{X} .*

1. If \mathcal{F} is almost bracket-generating, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2) such that

$$\mathcal{F} = \frac{\partial}{\partial x^1} + \text{span} \left(\frac{\partial}{\partial x^2} \right).$$

2. If \mathcal{F} is bracket-generating, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2) such that

$$\mathcal{F} = x^2 \left(\frac{\partial}{\partial x^1} + J \frac{\partial}{\partial x^2} \right) + \text{span} \left(\frac{\partial}{\partial x^2} \right),$$

where J is an arbitrary function on \mathcal{X} .

Proof. We will employ Cartan's method of equivalence in this proof. Let \mathcal{F} be a rank 1 point-affine distribution of constant type on a 2-dimensional manifold \mathcal{X} . A local framing – i.e., a pair of everywhere linearly independent vector fields (v_1, v_2) on \mathcal{X} – will be called *admissible* if v_1 is the distinguished vector field and

$$\mathcal{F} = v_1 + \text{span}(v_2).$$

If (v_1, v_2) is any admissible framing on \mathcal{X} , then any other admissible framing $(\tilde{v}_1, \tilde{v}_2)$ has the form

$$\tilde{v}_1 = v_1, \quad \tilde{v}_2 = b_2 v_2,$$

for some nonvanishing function b_2 on \mathcal{X} .

A local coframing $(\bar{\eta}^1, \bar{\eta}^2)$ on \mathcal{X} will be called *0-adapted* if it is the dual coframing of some admissible framing. Any two 0-adapted coframings on \mathcal{X} vary by a transformation of the form

$$\begin{bmatrix} \tilde{\eta}^1 \\ \tilde{\eta}^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & b_2 \end{bmatrix}^{-1} \begin{bmatrix} \bar{\eta}^1 \\ \bar{\eta}^2 \end{bmatrix}, \quad (3.1)$$

with $b_2 \neq 0$.

The 0-adapted coframings are the local sections of a principal fiber bundle $\pi : \mathcal{B}_0 \rightarrow \mathcal{X}$, with structure group $G_0 \subset GL(2)$ consisting of all invertible matrices of the form in (3.1). The right action of G_0 on sections $\sigma : \mathcal{X} \rightarrow \mathcal{B}_0$ is given by $\sigma \cdot g = g^{-1}\sigma$. (This is the reason for the inverse occurring in (3.1).)

There exist canonical 1-forms η^1, η^2 on \mathcal{B}_0 with the *reproducing property* that for any coframing $(\bar{\eta}^1, \bar{\eta}^2)$ given by a local section $\sigma : \mathcal{X} \rightarrow \mathcal{B}_0$,

$$\sigma^*(\eta^i) = \bar{\eta}^i.$$

These are referred to as the *semi-basic* forms on \mathcal{B}_0 . A standard argument shows that there also exists a (non-unique) 1-form β_2 (referred to as a *pseudo-connection form* or, more succinctly, a *connection form*), linearly independent from the semi-basic forms, and a function T_{12}^1 on \mathcal{B}_0 (referred to as a *torsion function*) such that

$$\begin{bmatrix} d\eta^1 \\ d\eta^2 \end{bmatrix} = - \begin{bmatrix} 0 & 0 \\ 0 & \beta_2 \end{bmatrix} \begin{bmatrix} \eta^1 \\ \eta^2 \end{bmatrix} + \begin{bmatrix} T_{12}^1 \eta^1 \wedge \eta^2 \\ 0 \end{bmatrix}. \quad (3.2)$$

(See [15] for details about the method of equivalence.) These are the *structure equations* of the G_0 -structure \mathcal{B}_0 . The semi-basic forms and the connection form together form a local coframing on \mathcal{B}_0 .

We proceed with the method of equivalence by examining how the function T_{12}^1 varies if we change from one 0-adapted coframing to another. A straightforward computation shows that under a transformation of the form (3.1), we have

$$\tilde{T}_{12}^1 = b_2 T_{12}^1. \quad (3.3)$$

Thus T_{12}^1 is a *relative invariant*: if it vanishes for any coframing at a point $x \in \mathcal{X}$, then it vanishes for every coframing at x .

At this point, we need to divide into cases based on whether or not T_{12}^1 vanishes.

Case 1. Suppose that $T_{12}^1 = 0$. Then the structure equations (3.2) contain no local invariants, so no further adaptations can be made. The tableau consisting of all matrices of the form in (3.2) is involutive with Cartan character $s_1 = 1$.

Given any 0-adapted coframing, consider the dual framing (v_1, v_2) . The structure equations imply that

$$[v_1, v_2] \equiv 0 \pmod{v_2};$$

therefore, \mathcal{F} is almost bracket-generating. In order to recover the normal form of Theorem 3.1, let $(\bar{\eta}^1, \bar{\eta}^2)$ be any 0-adapted local coframing on \mathcal{X} given by a local section $\sigma : \mathcal{X} \rightarrow \mathcal{B}_0$, and consider the pullbacks of the structure equations (3.2) via σ :

$$d\bar{\eta}^1 = 0, \quad d\bar{\eta}^2 = -\bar{\beta}_2 \wedge \bar{\eta}^2.$$

(Note that the pullback $\bar{\beta}_2$ of the connection form β_2 is a 1-form on \mathcal{X} , so it is no longer linearly independent from $\bar{\eta}^1, \bar{\eta}^2$.)

Since $\bar{\eta}^1$ is an exact 1-form, we can choose a local coordinate x^1 on \mathcal{X} such that $\bar{\eta}^1 = dx^1$. Then, since $\bar{\eta}^2$ is an integrable 1-form, we can choose a local coordinate x^2 on \mathcal{X} , functionally independent from x^1 , such that

$$\bar{\eta}^2 = \lambda dx^2$$

for some nonvanishing function λ on \mathcal{X} . By replacing the given coframing with the (also 0-adapted) coframing $(\bar{\eta}^1, \lambda^{-1}\bar{\eta}^2)$, we can assume that $\lambda \equiv 1$. Our 1-adapted coframing now has the form

$$\bar{\eta}^1 = dx^1, \quad \bar{\eta}^2 = dx^2.$$

The dual framing is

$$v_1 = \frac{\partial}{\partial x^1}, \quad v_2 = \frac{\partial}{\partial x^2},$$

and

$$\mathcal{F} = \frac{\partial}{\partial x^1} + \text{span} \left(\frac{\partial}{\partial x^2} \right).$$

Case 2. Suppose that $T_{12}^1 \neq 0$. (3.3) implies that, given any 0-adapted coframing $(\bar{\eta}^1, \bar{\eta}^2)$, we can perform a transformation of the form (3.1) to arrive at a new 0-adapted coframing for which $T_{12}^1 = 1$. Such a coframing will be called *1-adapted*.

Any two 1-adapted coframings on \mathcal{X} vary by a transformation of the form

$$\begin{bmatrix} \tilde{\eta}^1 \\ \tilde{\eta}^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} \bar{\eta}^1 \\ \bar{\eta}^2 \end{bmatrix}.$$

In other words, the 1-adapted coframings are the local sections of an (e) -structure $\mathcal{B}_1 \subset \mathcal{B}_0$. The projection $\pi : \mathcal{B}_1 \rightarrow \mathcal{X}$ is a diffeomorphism, and there is a *unique* 1-adapted coframing $(\bar{\eta}^1, \bar{\eta}^2)$ on \mathcal{X} . This coframing has structure equations

$$\begin{aligned} d\bar{\eta}^1 &= \bar{\eta}^1 \wedge \bar{\eta}^2, \\ d\bar{\eta}^2 &= T_{12}^2 \bar{\eta}^1 \wedge \bar{\eta}^2, \end{aligned} \tag{3.4}$$

for some function T_{12}^2 on \mathcal{X} .

Consider the dual framing (v_1, v_2) of a 1-adapted coframing. The structure equations (3.4) imply that

$$[v_1, v_2] \equiv -v_1 \pmod{v_2};$$

therefore, \mathcal{F} is bracket-generating. Since $\bar{\eta}^1$ is integrable but not exact, we can choose local coordinates x^1, x^2 on \mathcal{X} such that $\bar{\eta}^1 = \frac{1}{x^2} dx^1$. Then the first equation in (3.4) implies that

$$\bar{\eta}^2 = \frac{1}{x^2} (dx^2 - J dx^1)$$

for some function J on \mathcal{X} . The second equation then implies that

$$T_{12}^2 = x^2 J_{x^2} - J.$$

In terms of these local coordinates, the dual framing is

$$v_1 = x^2 \left(\frac{\partial}{\partial x^1} + J \frac{\partial}{\partial x^2} \right), \quad v_2 = x^2 \frac{\partial}{\partial x^2},$$

and since $\text{span}(v_2) = \text{span}\left(\frac{\partial}{\partial x^2}\right)$,

$$\mathcal{F} = x^2 \left(\frac{\partial}{\partial x^1} + J \frac{\partial}{\partial x^2} \right) + \text{span} \left(\frac{\partial}{\partial x^2} \right). \quad (3.5)$$

■

Remark 3.2. This choice of coordinates x^1, x^2 such that $\bar{\eta}^1 = \frac{1}{x^2} dx^1$ is unique up to coordinate transformations of the form

$$x^1 \rightarrow f(\tilde{x}^1), \quad x^2 \rightarrow f'(\tilde{x}^1)\tilde{x}^2. \quad (3.6)$$

Under a transformation of the form (3.6), we have

$$J(x^1, x^2) \rightarrow J(f(\tilde{x}^1), f'(\tilde{x}^1)\tilde{x}^2) - \frac{f''(\tilde{x}^1)}{f'(\tilde{x}^1)}\tilde{x}^2.$$

Therefore, an admissible change of coordinates can only change J by a linear function of x^2 , and point-affine distributions of this type locally depend essentially on one arbitrary function of 2 variables. In particular, the point-affine distribution (3.5) is locally equivalent to the “flat” case $J = 0$ if and only if $J = g(x^1)x^2$ for some function $g(x^1)$.

3.2 Equivalence problem for rank $(n - 1)$ point-affine distributions on n -manifolds

Theorem 3.1 is a special case of the following more general theorem:

Theorem 3.3. *Let \mathcal{F} be a rank $(n - 1)$ strictly affine point-affine distribution of constant type on an n -dimensional manifold \mathcal{X} . Let $L_{\mathcal{F}}$ be the associated direction distribution of rank $(n - 1)$, let v_1 denote the distinguished vector field, and let $\bar{\eta}^1$ be the unique 1-form on \mathcal{X} satisfying*

$$L_{\mathcal{F}} = (\bar{\eta}^1)^\perp, \quad \bar{\eta}^1(v_1) = 1.$$

Let k denote the Pfaff rank of $\bar{\eta}^1$, i.e., the unique integer such that

$$\bar{\eta}^1 \wedge (d\bar{\eta}^1)^k \neq 0, \quad \bar{\eta}^1 \wedge (d\bar{\eta}^1)^{k+1} = 0.$$

1. *If $(d\bar{\eta}^1)^{k+1} = 0$, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$ on which k is constant, there exist local coordinates (x^1, \dots, x^n) such that*

$$\begin{aligned} \mathcal{F} = & \left(1 + \sum_{r=1}^k x^{k+r+1} J_{k+r+1} \right) \frac{\partial}{\partial x^1} + \sum_{r=1}^k \left(J_{k+r+1} \frac{\partial}{\partial x^{r+1}} - J_{r+1} \frac{\partial}{\partial x^{k+r+1}} \right) \\ & + \text{span} \left(\left(\frac{\partial}{\partial x^2} + x^{k+2} \frac{\partial}{\partial x^1} \right), \dots, \left(\frac{\partial}{\partial x^{k+1}} + x^{2k+1} \frac{\partial}{\partial x^1} \right), \frac{\partial}{\partial x^{k+2}}, \dots, \frac{\partial}{\partial x^n} \right), \end{aligned}$$

where J_2, \dots, J_{2k+1} are arbitrary functions on \mathcal{X} .

2. If $(d\bar{\eta}^1)^{k+1} \neq 0$, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$ on which k is constant, there exist local coordinates (x^1, \dots, x^n) such that

$$\begin{aligned} \mathcal{F} = & \left(x^{2k+2} + \sum_{r=1}^k x^{k+r+1} J_{k+r+1} \right) \frac{\partial}{\partial x^1} + \sum_{r=1}^k \left(J_{k+r+1} \frac{\partial}{\partial x^{r+1}} - J_{r+1} \frac{\partial}{\partial x^{k+r+1}} \right) - J_1 \frac{\partial}{\partial x^{2k+2}} \\ & + \text{span} \left(\left(x^{2k+2} \left(\frac{\partial}{\partial x^2} + x^{k+2} \frac{\partial}{\partial x^1} \right) - J_2 \frac{\partial}{\partial x^{2k+2}} \right), \dots, \right. \\ & \quad \left(x^{2k+2} \left(\frac{\partial}{\partial x^{k+1}} + x^{2k+1} \frac{\partial}{\partial x^1} \right) - J_{k+1} \frac{\partial}{\partial x^{2k+2}} \right), \\ & \quad \left(\frac{\partial}{\partial x^{k+2}} - J_{k+2} \frac{\partial}{\partial x^{2k+2}} \right), \dots, \left(\frac{\partial}{\partial x^{2k+1}} - J_{2k+1} \frac{\partial}{\partial x^{2k+2}} \right), \\ & \quad \left. \frac{\partial}{\partial x^{2k+2}}, \dots, \frac{\partial}{\partial x^n} \right), \end{aligned}$$

where J_1, \dots, J_{2k+1} are arbitrary functions on \mathcal{X} .

Proof. Let \mathcal{F} be a rank $(n-1)$ point-affine distribution of constant type on an n -dimensional manifold \mathcal{X} . A local framing (v_1, \dots, v_n) on \mathcal{X} will be called *admissible* if v_1 is the distinguished vector field and

$$\mathcal{F} = v_1 + \text{span}(v_2, \dots, v_n).$$

A local coframing $(\bar{\eta}^1, \dots, \bar{\eta}^n)$ on \mathcal{X} will be called *admissible* if, for any admissible framing (v_1, \dots, v_n) ,

$$\begin{aligned} \bar{\eta}^1(v_1) &= 1, \\ \bar{\eta}^1(v_j) &= 0, \quad 2 \leq j \leq n, \\ \bar{\eta}^j(v_1) &= 0, \quad 2 \leq j \leq n. \end{aligned}$$

Any two admissible coframings on \mathcal{X} vary by a transformation of the form

$$\begin{bmatrix} \tilde{\eta}^1 \\ \tilde{\eta}^2 \\ \vdots \\ \tilde{\eta}^n \end{bmatrix} = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & & & \\ \vdots & & A & \\ 0 & & & \end{bmatrix}^{-1} \begin{bmatrix} \bar{\eta}^1 \\ \bar{\eta}^2 \\ \vdots \\ \bar{\eta}^n \end{bmatrix}$$

for some matrix $A \in GL(n-1, \mathbb{R})$.

An admissible coframing will be called *0-adapted* if

$$d\bar{\eta}^1 \equiv \sum_{r=1}^k \bar{\eta}^{r+1} \wedge \bar{\eta}^{k+r+1} \pmod{\bar{\eta}^1},$$

where k is the Pfaff rank of $\bar{\eta}^1$. Since the Cartan system

$$\mathcal{C}(\bar{\eta}^1) = \{\bar{\eta}^1, \dots, \bar{\eta}^{2k+1}\}$$

and the system

$$v_1^\perp = \{\bar{\eta}^2, \dots, \bar{\eta}^n\}$$

are both well-defined independent of the choice of 0-adapted coframing, the subsystem

$$\mathcal{C}(\bar{\eta}^1) \cap v_1^\perp = \{\bar{\eta}^2, \dots, \bar{\eta}^{2k+1}\}$$

is also well-defined. Thus any two 0-adapted coframings on \mathcal{X} vary by a transformation of the form

$$\begin{bmatrix} \tilde{\eta}^1 \\ \tilde{\eta}^2 \\ \vdots \\ \tilde{\eta}^{2k+1} \\ \tilde{\eta}^{2k+2} \\ \vdots \\ \tilde{\eta}^n \end{bmatrix} = \begin{bmatrix} 1 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & & & & 0 & \cdots & 0 \\ \vdots & & A & & \vdots & & \vdots \\ 0 & & & & 0 & \cdots & 0 \\ 0 & & & & & & \\ \vdots & & B & & C & & \\ 0 & & & & & & \end{bmatrix}^{-1} \begin{bmatrix} \bar{\eta}^1 \\ \bar{\eta}^2 \\ \vdots \\ \bar{\eta}^{2k+1} \\ \bar{\eta}^{2k+2} \\ \vdots \\ \bar{\eta}^n \end{bmatrix}, \quad (3.7)$$

where $A \in Sp(2k, \mathbb{R})$, $B \in M_{(n-2k-1) \times (2k)}(\mathbb{R})$, $C \in GL(n-2k-1, \mathbb{R})$.

The bundle $\pi : \mathcal{B}_0 \rightarrow \mathcal{X}$ of 0-adapted coframings with structure group G_0 consisting of matrices of the form (3.7) is defined as in Section 3.1, as are the semi-basic forms η^1, \dots, η^n on \mathcal{B}_0 . The first structure equation of \mathcal{B}_0 is

$$d\eta^1 = \eta^1 \wedge \phi + \sum_{r=1}^k \eta^{r+1} \wedge \eta^{k+r+1},$$

where $\phi \in \pi^*(v_1^\perp)$. There are two cases to consider:

1. $\phi \in \mathcal{C}(\eta^1)$;
2. $\phi \notin \mathcal{C}(\eta^1)$.

Case 1: $\phi \in \mathcal{C}(\eta^1)$. Then

$$(d\eta^1)^{k+1} = 0.$$

Let $\sigma : \mathcal{X} \rightarrow \mathcal{B}_0$ be any local section of \mathcal{B}_0 (i.e., a 0-adapted coframing $(\bar{\eta}^1, \dots, \bar{\eta}^n)$ on \mathcal{X}). By the reproducing property of the semi-basic forms,

$$(d\bar{\eta}^1)^{k+1} = 0,$$

so by the Pfaff theorem there exist local coordinates x^1, \dots, x^{2k+1} on \mathcal{X} such that

$$\bar{\eta}^1 = dx^1 - \sum_{r=1}^k x^{k+r+1} dx^{r+1}. \quad (3.8)$$

We can write

$$\bar{\phi} = \sum_{j=1}^{2k} J_{j+1} \bar{\eta}^{j+1}$$

for some functions J_2, \dots, J_{2k+1} on \mathcal{X} . Then we have

$$\begin{aligned} d\bar{\eta}^1 &= \bar{\eta}^1 \wedge \left(\sum_{j=1}^{2k} J_{j+1} \bar{\eta}^{j+1} \right) + \sum_{r=1}^k \bar{\eta}^{r+1} \wedge \bar{\eta}^{k+r+1} \\ &= \sum_{r=1}^k (\bar{\eta}^{r+1} + J_{k+r+1} \bar{\eta}^1) \wedge (\bar{\eta}^{k+r+1} - J_{r+1} \bar{\eta}^1) = \sum_{r=1}^k dx^{r+1} \wedge dx^{k+r+1}. \end{aligned}$$

Thus we can choose a 0-adapted coframing with

$$\left. \begin{aligned} \bar{\eta}^{s+1} &= dx^{s+1} - J_{k+s+1} \bar{\eta}^1 \\ \bar{\eta}^{k+s+1} &= dx^{k+s+1} + J_{s+1} \bar{\eta}^1 \end{aligned} \right\}, \quad 1 \leq s \leq k. \quad (3.9)$$

For reasons of dimension, the system

$$v_1^\perp = \{\bar{\eta}^2, \dots, \bar{\eta}^n\}$$

is Frobenius; therefore, we can extend the local coordinates x^1, \dots, x^{2k+1} to a full local coordinate system x^1, \dots, x^n with the property that

$$v_1^\perp = \{\bar{\eta}^2, \dots, \bar{\eta}^{2k+1}, dx^{2k+2}, \dots, dx^n\};$$

we can then choose the remainder of our 0-adapted coframing to be

$$\bar{\eta}^j = dx^j, \quad 2k+2 \leq j \leq n. \quad (3.10)$$

The dual framing to the coframing defined by (3.8), (3.9), and (3.10) is:

$$\begin{aligned} v_1 &= \left(1 + \sum_{r=1}^k x^{k+r+1} J_{k+r+1} \right) \frac{\partial}{\partial x^1} + \sum_{r=1}^k \left(J_{k+r+1} \frac{\partial}{\partial x^{r+1}} - J_{r+1} \frac{\partial}{\partial x^{k+r+1}} \right), \\ v_s &= \frac{\partial}{\partial x^s} + x^{k+s} \frac{\partial}{\partial x^1}, \quad 2 \leq s \leq k+1, \\ v_j &= \frac{\partial}{\partial x^j}, \quad k+2 \leq j \leq n. \end{aligned}$$

The corresponding point-affine distribution \mathcal{F} is:

$$\begin{aligned} \mathcal{F} &= \left(1 + \sum_{r=1}^k x^{k+r+1} J_{k+r+1} \right) \frac{\partial}{\partial x^1} + \sum_{r=1}^k \left(J_{k+r+1} \frac{\partial}{\partial x^{r+1}} - J_{r+1} \frac{\partial}{\partial x^{k+r+1}} \right) \\ &\quad + \text{span} \left(\left(\frac{\partial}{\partial x^2} + x^{k+2} \frac{\partial}{\partial x^1} \right), \dots, \left(\frac{\partial}{\partial x^{k+1}} + x^{2k+1} \frac{\partial}{\partial x^1} \right), \frac{\partial}{\partial x^{k+2}}, \dots, \frac{\partial}{\partial x^n} \right). \end{aligned}$$

Note that our choices of local coordinates were arbitrary up to at most functions of $(n-1)$ variables, while the local invariants J_2, \dots, J_{2k+1} are arbitrary functions of n variables. Therefore, point-affine distributions of this type locally depend essentially on $2k$ arbitrary functions of n variables.

Case 2: $\phi \notin \mathcal{C}(\eta^1)$. Then

$$(d\eta^1)^{k+1} \neq 0.$$

Let $\sigma : \mathcal{X} \rightarrow \mathcal{B}_0$ be any local section of \mathcal{B}_0 (i.e., a 0-adapted coframing $(\bar{\eta}^1, \dots, \bar{\eta}^n)$ on \mathcal{X}). Because $d\eta^1$ is semi-basic, it follows from the reproducing property of the semi-basic forms that

$$(d\bar{\eta}^1)^{k+1} \neq 0.$$

By the Pfaff theorem, there exist local coordinates x^1, \dots, x^{2k+2} on \mathcal{X} such that

$$\bar{\eta}^1 = \frac{1}{x^{2k+2}} \left(dx^1 - \sum_{r=1}^k x^{k+r+1} dx^{r+1} \right). \quad (3.11)$$

Then we have

$$\begin{aligned} d\bar{\eta}^1 &= \bar{\eta}^1 \wedge \bar{\phi} + \sum_{r=1}^k \bar{\eta}^{r+1} \wedge \bar{\eta}^{k+r+1} \\ &= \bar{\eta}^1 \wedge \left(\frac{dx^{2k+2}}{x^{2k+2}} \right) + \frac{1}{x^{2k+2}} \sum_{r=1}^k dx^{r+1} \wedge dx^{k+r+1}. \end{aligned}$$

Therefore,

$$\sum_{r=1}^k \bar{\eta}^{r+1} \wedge \bar{\eta}^{k+r+1} \equiv \frac{1}{x^{2k+2}} \sum_{r=1}^k dx^{r+1} \wedge dx^{k+r+1} \pmod{\bar{\eta}^1}.$$

Thus we can choose a 0-adapted coframing with

$$\left. \begin{aligned} \bar{\eta}^{s+1} &= \frac{1}{x^{2k+2}} (dx^{s+1} - J_{k+s+1} \bar{\eta}^1) \\ \bar{\eta}^{k+s+1} &= dx^{k+s+1} + J_{s+1} \bar{\eta}^1 \end{aligned} \right\}, \quad 1 \leq s \leq k, \quad (3.12)$$

for some functions J_2, \dots, J_{2k+1} on \mathcal{X} . Then we have

$$\bar{\phi} \equiv \frac{1}{x^{2k+2}} \left(dx^{2k+2} + \sum_{j=1}^{2k} J_{j+1} \bar{\eta}^{j+1} \right) \pmod{\bar{\eta}^1}.$$

The requirement that $\phi \in \pi^*(v_1^\perp)$ is equivalent to $\bar{\phi} \in v_1^\perp$, and this determines a unique function J_1 on \mathcal{X} such that

$$\bar{\phi} = \frac{1}{x^{2k+2}} \left(dx^{2k+2} + \sum_{j=0}^{2k} J_{j+1} \bar{\eta}^{j+1} \right).$$

We can now set

$$\bar{\eta}^{2k+2} = \bar{\phi} = \frac{1}{x^{2k+2}} \left(dx^{2k+2} + \sum_{j=0}^{2k} J_{j+1} \bar{\eta}^{j+1} \right), \quad (3.13)$$

and a similar argument to that given in the previous case shows that we can extend the local coordinates x^1, \dots, x^{2k+2} to a full local coordinate system x^1, \dots, x^n and choose the remainder of our 0-adapted coframing to be

$$\bar{\eta}^j = dx^j, \quad 2k+3 \leq j \leq n. \quad (3.14)$$

The dual framing to the coframing defined by (3.11), (3.12), (3.13), and (3.14) is:

$$\begin{aligned} v_1 &= \left(x^{2k+2} + \sum_{r=1}^k x^{k+r+1} J_{k+r+1} \right) \frac{\partial}{\partial x^1} + \sum_{r=1}^k \left(J_{k+r+1} \frac{\partial}{\partial x^{r+1}} - J_{r+1} \frac{\partial}{\partial x^{k+r+1}} \right) - J_1 \frac{\partial}{\partial x^{2k+2}}, \\ v_s &= x^{2k+2} \left(\frac{\partial}{\partial x^s} + x^{k+s} \frac{\partial}{\partial x^1} \right) - J_s \frac{\partial}{\partial x^{2k+2}}, \quad 2 \leq s \leq k+1, \\ v_s &= \frac{\partial}{\partial x^s} - J_s \frac{\partial}{\partial x^{2k+2}}, \quad k+2 \leq s \leq 2k+1, \\ v_{2k+2} &= x^{2k+2} \frac{\partial}{\partial x^{2k+2}}, \quad v_j = \frac{\partial}{\partial x^j}, \quad 2k+3 \leq j \leq n. \end{aligned}$$

Since $\text{span}(v_{2k+2}) = \text{span}\left(\frac{\partial}{\partial x^{2k+2}}\right)$, the corresponding point-affine distribution \mathcal{F} is:

$$\begin{aligned} \mathcal{F} = & \left(x^{2k+2} + \sum_{r=1}^k x^{k+r+1} J_{k+r+1} \right) \frac{\partial}{\partial x^1} + \sum_{r=1}^k \left(J_{k+r+1} \frac{\partial}{\partial x^{r+1}} - J_{r+1} \frac{\partial}{\partial x^{k+r+1}} \right) - J_1 \frac{\partial}{\partial x^{2k+2}} \\ & + \text{span} \left(\left(x^{2k+2} \left(\frac{\partial}{\partial x^2} + x^{k+2} \frac{\partial}{\partial x^1} \right) - J_2 \frac{\partial}{\partial x^{2k+2}} \right), \dots, \right. \\ & \left. \left(x^{2k+2} \left(\frac{\partial}{\partial x^{k+1}} + x^{2k+1} \frac{\partial}{\partial x^1} \right) - J_{k+1} \frac{\partial}{\partial x^{2k+2}} \right), \right. \\ & \left. \left(\frac{\partial}{\partial x^{k+2}} - J_{k+2} \frac{\partial}{\partial x^{2k+2}} \right), \dots, \left(\frac{\partial}{\partial x^{2k+1}} - J_{2k+1} \frac{\partial}{\partial x^{2k+2}} \right), \frac{\partial}{\partial x^{2k+2}}, \dots, \frac{\partial}{\partial x^n} \right). \end{aligned}$$

Note that our choices of local coordinates were arbitrary up to at most functions of $(n-1)$ variables, while the local invariants J_1, \dots, J_{2k+1} are arbitrary functions of n variables. Therefore, point-affine distributions of this type locally depend essentially on $2k+1$ arbitrary functions of n variables. \blacksquare

Corollary 1. *Let \mathcal{F} be a rank 2 strictly affine point-affine distribution of constant type on a 3-dimensional manifold \mathcal{X} .*

1. *If \mathcal{F} is almost bracket-generating, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3) such that*

$$\mathcal{F} = \frac{\partial}{\partial x^1} + \text{span} \left(\frac{\partial}{\partial x^2}, \frac{\partial}{\partial x^3} \right).$$

2. *If \mathcal{F} is bracket-generating and $L_{\mathcal{F}}$ is Frobenius, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3) such that*

$$\mathcal{F} = \left(x^2 \frac{\partial}{\partial x^1} - J_1 \frac{\partial}{\partial x^2} \right) + \text{span} \left(\frac{\partial}{\partial x^2}, \frac{\partial}{\partial x^3} \right),$$

where J_1 is an arbitrary function on \mathcal{X} .

3. *If \mathcal{F} is bracket-generating and $L_{\mathcal{F}}$ is a contact distribution, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3) such that*

$$\mathcal{F} = \left((1 + x^3 J_3) \frac{\partial}{\partial x^1} + J_3 \frac{\partial}{\partial x^2} - J_2 \frac{\partial}{\partial x^3} \right) + \text{span} \left(x^3 \frac{\partial}{\partial x^1} + \frac{\partial}{\partial x^2}, \frac{\partial}{\partial x^3} \right),$$

where J_2, J_3 are arbitrary functions on \mathcal{X} .

Proof. Let $\bar{\eta}^1$ be as in Theorem 3.3.

1. If \mathcal{F} is almost bracket-generating, then $d\bar{\eta}^1 = 0$, so $k = 0$ and Case 1 of Theorem 3.3 applies.

2. If \mathcal{F} is bracket-generating and $L_{\mathcal{F}}$ is Frobenius, then

$$\bar{\eta}^1 \wedge d\bar{\eta}^1 = 0, \quad d\bar{\eta}^1 \neq 0,$$

so $k = 0$ and Case 2 of Theorem 3.3 applies.

3. If \mathcal{F} is bracket-generating and $L_{\mathcal{F}}$ is a contact distribution, then

$$\bar{\eta}^1 \wedge d\bar{\eta}^1 \neq 0, \quad (d\bar{\eta}^1)^2 = 0,$$

so $k = 1$ and Case 1 of Theorem 3.3 applies. \blacksquare

3.3 Equivalence problem for rank 1 point-affine distributions on 3-manifolds

In this section we consider point-affine equivalence for rank 1 point-affine distributions \mathcal{F} on 3-dimensional manifolds. We will assume that \mathcal{F} is bracket-generating or almost bracket-generating, since otherwise the equivalence problem may be reduced to a problem on a 2-dimensional manifold.

Theorem 3.4. *Let \mathcal{F} be a rank 1 strictly affine point-affine distribution of constant type on a 3-dimensional manifold \mathcal{X} .*

1. *If \mathcal{F} is almost bracket-generating, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3) such that*

$$\mathcal{F} = \left(\frac{\partial}{\partial x^1} + x^3 \frac{\partial}{\partial x^2} + J \frac{\partial}{\partial x^3} \right) + \text{span} \left(\frac{\partial}{\partial x^3} \right),$$

where J is an arbitrary function on \mathcal{X} .

2. *If \mathcal{F} is bracket-generating and $L_{\mathcal{F}^2}$ (which must have rank 2) is Frobenius, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3) such that*

$$\mathcal{F} = \left(x^2 \frac{\partial}{\partial x^1} + x^3 \frac{\partial}{\partial x^2} + J \frac{\partial}{\partial x^3} \right) + \text{span} \left(\frac{\partial}{\partial x^3} \right),$$

where J is an arbitrary function on \mathcal{X} .

3. *If \mathcal{F} is bracket-generating and $L_{\mathcal{F}^2}$ is not Frobenius, then in a sufficiently small neighborhood of any point $x \in \mathcal{X}$, there exist local coordinates (x^1, x^2, x^3) such that*

$$\mathcal{F} = \left(\frac{\partial}{\partial x^1} + J \left(x^3 \frac{\partial}{\partial x^1} + \frac{\partial}{\partial x^2} + H \frac{\partial}{\partial x^3} \right) \right) + \text{span} \left(x^3 \frac{\partial}{\partial x^1} + \frac{\partial}{\partial x^2} + H \frac{\partial}{\partial x^3} \right),$$

where H, J are arbitrary functions on \mathcal{X} satisfying $\frac{\partial H}{\partial x^1} \neq 0$.

Compare Theorem 3.4 with Theorem 2.6; in each case there is a new invariant J which reflects the restriction that v_1 is a fixed vector field in \mathcal{F} .

Proof. A local framing of (v_1, v_2, v_3) on \mathcal{X} will be called *admissible* if v_1 is the distinguished vector field and

$$\mathcal{F} = v_1 + \text{span}(v_2).$$

If (v_1, v_2, v_3) is an admissible framing on \mathcal{X} , then any other admissible framing $(\tilde{v}_1, \tilde{v}_2, \tilde{v}_3)$ has the form

$$\tilde{v}_1 = v_1, \quad \tilde{v}_2 = b_2 v_2, \quad \tilde{v}_3 = a_3 v_1 + b_3 v_2 + c_3 v_3,$$

for some functions a_3, b_2, b_3, c_3 on \mathcal{X} which satisfy $b_2 c_3 \neq 0$.

A local coframing $(\bar{\eta}^1, \bar{\eta}^2, \bar{\eta}^3)$ on \mathcal{X} will be called *0-adapted* if it is the dual coframing of some admissible framing. Any two 0-adapted coframings on \mathcal{X} vary by a transformation of the form

$$\begin{bmatrix} \tilde{\eta}^1 \\ \tilde{\eta}^2 \\ \tilde{\eta}^3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & a_3 \\ 0 & b_2 & b_3 \\ 0 & 0 & c_3 \end{bmatrix}^{-1} \begin{bmatrix} \bar{\eta}^1 \\ \bar{\eta}^2 \\ \bar{\eta}^3 \end{bmatrix}, \quad (3.15)$$

with $b_2 c_3 \neq 0$.

The 0-adapted coframings are the local sections of a principal fiber bundle $\mathcal{B}_0 \rightarrow \mathcal{X}$, with structure group $G_0 \subset GL(3)$ consisting of all invertible matrices of the form in (3.15). The structure equations on \mathcal{B}_0 take the form

$$\begin{bmatrix} d\eta^1 \\ d\eta^2 \\ d\eta^3 \end{bmatrix} = - \begin{bmatrix} 0 & 0 & \alpha_3 \\ 0 & \beta_2 & \beta_3 \\ 0 & 0 & \gamma_3 \end{bmatrix} \begin{bmatrix} \eta^1 \\ \eta^2 \\ \eta^3 \end{bmatrix} + \begin{bmatrix} T_{12}^1 \eta^1 \wedge \eta^2 \\ 0 \\ T_{12}^3 \eta^1 \wedge \eta^2 \end{bmatrix}.$$

Under a transformation of the form (3.15), the torsion functions T_{12}^1, T_{12}^3 transform as follows:

$$\tilde{T}_{12}^1 = b_2 T_{12}^1 - \frac{a_3 b_2}{c_3} T_{12}^3, \quad \tilde{T}_{12}^3 = \frac{b_2}{c_3} T_{12}^3. \quad (3.16)$$

The function T_{12}^3 is a relative invariant, and the assumption that \mathcal{F} is (almost) bracket-generating implies that $T_{12}^3 \neq 0$. So, given any 0-adapted coframing (η^1, η^2, η^3) , (3.16) implies that we can perform a transformation of the form (3.15) to arrive at a new 0-adapted coframing for which $T_{12}^3 = 1, T_{12}^1 = 0$. Such a coframing will be called *1-adapted*.

Any two 1-adapted coframings on \mathcal{X} vary by a transformation of the form

$$\begin{bmatrix} \tilde{\eta}^1 \\ \tilde{\eta}^2 \\ \tilde{\eta}^3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & b_2 & b_3 \\ 0 & 0 & b_2 \end{bmatrix}^{-1} \begin{bmatrix} \eta^1 \\ \eta^2 \\ \eta^3 \end{bmatrix}, \quad (3.17)$$

with $b_2 \neq 0$. The 1-adapted coframings are the local sections of a principal fiber bundle $\mathcal{B}_1 \subset \mathcal{B}_0$, with structure group G_1 consisting of all matrices of the form in (3.17). When restricted to \mathcal{B}_1 , the connection forms $\alpha_3, \gamma_3 - \beta_2$ become semi-basic, thereby introducing new torsion terms into the structure equations of \mathcal{B}_1 . By adding multiples of the semi-basic forms to the connection forms β_2, β_3 so as to absorb as much of the torsion as possible, we can arrange that the structure equations of \mathcal{B}_1 take the form

$$\begin{bmatrix} d\eta^1 \\ d\eta^2 \\ d\eta^3 \end{bmatrix} = - \begin{bmatrix} 0 & 0 & 0 \\ 0 & \beta_2 & \beta_3 \\ 0 & 0 & \beta_2 \end{bmatrix} \begin{bmatrix} \eta^1 \\ \eta^2 \\ \eta^3 \end{bmatrix} + \begin{bmatrix} (T_{13}^1 \eta^1 + T_{23}^1 \eta^2) \wedge \eta^3 \\ 0 \\ \eta^1 \wedge \eta^2 + T_{13}^3 \eta^1 \wedge \eta^3 \end{bmatrix}.$$

Under a transformation of the form (3.17), we have

$$\tilde{T}_{13}^1 = b_2 T_{13}^1, \quad \tilde{T}_{23}^1 = b_2^2 T_{23}^1, \quad \tilde{T}_{13}^3 = T_{13}^3 + \frac{b_3}{b_2}.$$

The functions T_{13}^1, T_{23}^1 are relative invariants; however, given any 1-adapted coframing (η^1, η^2, η^3) , we can perform a transformation of the form (3.17) to arrive at a new 1-adapted coframing for which $T_{13}^3 = 0$. Such a coframing will be called *2-adapted*.

Any two 2-adapted coframings on \mathcal{X} vary by a transformation of the form

$$\begin{bmatrix} \tilde{\eta}^1 \\ \tilde{\eta}^2 \\ \tilde{\eta}^3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & b_2 & 0 \\ 0 & 0 & b_2 \end{bmatrix}^{-1} \begin{bmatrix} \eta^1 \\ \eta^2 \\ \eta^3 \end{bmatrix}, \quad (3.18)$$

with $b_2 \neq 0$. The 2-adapted coframings are the local sections of a principal fiber bundle $\mathcal{B}_2 \subset \mathcal{B}_1$, with structure group G_2 consisting of all matrices of the form in (3.18). When restricted to \mathcal{B}_2 , the connection form β_3 becomes semi-basic, thereby introducing new torsion terms into the

structure equations of \mathcal{B}_2 . By adding multiples of the semi-basic forms to the connection form β_2 so as to absorb as much of the torsion as possible, we can arrange that the structure equations of \mathcal{B}_2 take the form

$$\begin{bmatrix} d\eta^1 \\ d\eta^2 \\ d\eta^3 \end{bmatrix} = - \begin{bmatrix} 0 & 0 & 0 \\ 0 & \beta_2 & 0 \\ 0 & 0 & \beta_2 \end{bmatrix} \begin{bmatrix} \eta^1 \\ \eta^2 \\ \eta^3 \end{bmatrix} + \begin{bmatrix} (T_{13}^1\eta^1 + T_{23}^1\eta^2) \wedge \eta^3 \\ T_{13}^2\eta^1 \wedge \eta^3 \\ \eta^1 \wedge \eta^2 \end{bmatrix}. \quad (3.19)$$

Under a transformation of the form (3.18), we have

$$\tilde{T}_{13}^1 = b_2 T_{13}^1, \quad \tilde{T}_{23}^1 = b_2^2 T_{23}^1, \quad \tilde{T}_{13}^2 = T_{13}^2. \quad (3.20)$$

Thus the functions T_{13}^1, T_{23}^1 are relative invariants, while T_{13}^2 is a well-defined function on \mathcal{X} . As in Section 3.1, we will divide into cases based on the vanishing/nonvanishing of T_{23}^1 and T_{13}^1 .

Case 1. Suppose that $T_{23}^1 = T_{13}^1 = 0$. Since T_{13}^2 is well-defined on \mathcal{X} , no further adaptations can be made based on the structure equations (3.19). The tableau consisting of all matrices of the form in (3.19) is not involutive, so in principle the next step in the method of equivalence would be to prolong the structure equations. However, equations (3.19) already contain enough information to produce a local normal form for such structures, so we will proceed in this direction.

Given any 1-adapted coframing, consider the dual framing (v_1, v_2, v_3) . The structure equations (3.19) imply that

$$[v_1, v_2] \equiv -v_3 \pmod{v_2}, \quad [v_2, v_3] \equiv 0 \pmod{v_2, v_3}, \quad [v_1, v_3] \equiv 0 \pmod{v_2, v_3}.$$

The first equation implies that $L_{\mathcal{F}^2}$ is spanned by $\{v_2, v_3\}$; the second implies that $L_{\mathcal{F}^2}$ is a Frobenius distribution, and the third implies that \mathcal{F} is almost bracket-generating. Thus, this case corresponds to Case 1 of Theorem 3.4.

In order to recover the normal form of Theorem 3.4, let $(\bar{\eta}^1, \bar{\eta}^2, \bar{\eta}^3)$ be any 1-adapted local coframing on \mathcal{X} given by a local section $\sigma : \mathcal{X} \rightarrow \mathcal{B}_2$, and consider the pullbacks of the structure equations (3.19) via σ :

$$\begin{aligned} d\bar{\eta}^1 &= 0, \\ d\bar{\eta}^2 &= -\bar{\beta}_2 \wedge \bar{\eta}^2 + \bar{T}_{13}^2 \bar{\eta}^1 \wedge \bar{\eta}^3, \\ d\bar{\eta}^3 &= -\bar{\beta}_2 \wedge \bar{\eta}^3 + \bar{\eta}^1 \wedge \bar{\eta}^2. \end{aligned} \quad (3.21)$$

Since $\bar{\eta}^1$ is an exact 1-form, we can choose a local coordinate x^1 on \mathcal{X} such that $\bar{\eta}^1 = dx^1$. Then, since $\bar{\eta}^3$ is a contact form with

$$d\bar{\eta}^3 \equiv 0 \pmod{\{\bar{\eta}^3, dx^1\}},$$

there exist local coordinates x^2, x^3 , functionally independent from x^1 , and a nonvanishing function λ on \mathcal{X} such that

$$\bar{\eta}^3 = \lambda(dx^2 - x^3 dx^1).$$

By replacing the given coframing with the (also 1-adapted) coframing $(\bar{\eta}^1, \lambda^{-1}\bar{\eta}^2, \lambda^{-1}\bar{\eta}^3)$, we can assume that $\lambda \equiv 1$. The third equation in (3.21) then implies that

$$\bar{\eta}^2 = dx^3 + Bdx^1 + C(dx^2 - x^3 dx^1)$$

for some functions B, C on \mathcal{X} , and that

$$\bar{\beta}_2 = Cdx^1 + D(dx^2 - x^3 dx^1)$$

for some additional function D on \mathcal{X} . Finally, the second equation in (3.21) implies that

$$C = \frac{1}{2}B_{x^3}, \quad D = \frac{1}{2}B_{x^3x^3}.$$

Our 2-adapted coframing now has the form

$$\begin{aligned} \bar{\eta}^1 &= dx^1, \\ \bar{\eta}^2 &= dx^3 + B dx^1 + \frac{1}{2}B_{x^3}(dx^2 - x^3 dx^1), \\ \bar{\eta}^3 &= dx^2 - x^3 dx^1. \end{aligned} \tag{3.22}$$

The dual framing is

$$v_1 = \frac{\partial}{\partial x^1} + x^3 \frac{\partial}{\partial x^2} - B \frac{\partial}{\partial x^3}, \quad v_2 = \frac{\partial}{\partial x^3}, \quad v_3 = \frac{\partial}{\partial x^2} - \frac{1}{2}B_{x^3} \frac{\partial}{\partial x^3};$$

setting $J = -B$, we have:

$$\mathcal{F} = v_1 + \text{span}(v_2) = \left(\frac{\partial}{\partial x^1} + x^3 \frac{\partial}{\partial x^2} + J \frac{\partial}{\partial x^3} \right) + \text{span} \left(\frac{\partial}{\partial x^3} \right).$$

Comparing the structure equations (3.21) with those of the coframing (3.22), we see that

$$T_{13}^2 = J_{x^2} - \frac{1}{2}(J_{x^1x^3} + x^3 J_{x^2x^3} + J J_{x^3x^3}) + \frac{1}{4}(J_{x^3})^2.$$

Case 2. Suppose that $T_{23}^1 = 0$, $T_{13}^1 \neq 0$. (3.20) implies that, given any 2-adapted coframing $(\bar{\eta}^1, \bar{\eta}^2, \bar{\eta}^3)$, we can perform a transformation of the form (3.18) to arrive at a new 2-adapted coframing for which $T_{13}^1 = 1$. Such a coframing will be called *3-adapted*.

Any two 3-adapted coframings on \mathcal{X} vary by a transformation of the form

$$\begin{bmatrix} \tilde{\eta}^1 \\ \tilde{\eta}^2 \\ \tilde{\eta}^3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} \bar{\eta}^1 \\ \bar{\eta}^2 \\ \bar{\eta}^3 \end{bmatrix}.$$

In other words, the 3-adapted coframings are the local sections of an (e) -structure $\mathcal{B}_3 \subset \mathcal{B}_2$. The projection $\pi : \mathcal{B}_3 \rightarrow \mathcal{X}$ is a diffeomorphism, and there is a *unique* 3-adapted coframing $(\bar{\eta}^1, \bar{\eta}^2, \bar{\eta}^3)$ on \mathcal{X} .

When restricted to \mathcal{B}_3 , the connection form β_2 becomes semi-basic. The first structure equation in (3.19) now takes the form

$$d\eta^1 = \eta^1 \wedge \eta^3.$$

Differentiating this equation yields

$$\beta_2 \wedge \eta^1 \wedge \eta^3 = 0;$$

therefore, β_2 is a linear combination of η^1 and η^3 , and the structure equations of \mathcal{B}_3 take the form

$$\begin{aligned} d\eta^1 &= \eta^1 \wedge \eta^3, \\ d\eta^2 &= T_{12}^2 \eta^1 \wedge \eta^2 + T_{13}^2 \eta^1 \wedge \eta^3 + T_{23}^2 \eta^2 \wedge \eta^3, \\ d\eta^3 &= \eta^1 \wedge \eta^2 + T_{12}^2 \eta^1 \wedge \eta^3. \end{aligned} \tag{3.23}$$

Given any 3-adapted coframing, consider the dual framing (v_1, v_2, v_3) . The structure equations (3.23) imply that

$$[v_1, v_2] \equiv -v_3 \pmod{v_2}, \quad [v_2, v_3] \equiv 0 \pmod{v_2}, \quad [v_1, v_3] \equiv -v_1 \pmod{v_2}.$$

The first equation implies that $L_{\mathcal{F}^2}$ is spanned by $\{v_2, v_3\}$; the second implies that $L_{\mathcal{F}^2}$ is a Frobenius distribution, and the third implies that \mathcal{F} is bracket-generating. Thus, this case corresponds to Case 2 of Theorem 3.4.

In order to recover the normal form of Theorem 3.4, let $(\bar{\eta}^1, \bar{\eta}^2, \bar{\eta}^3)$ be the 3-adapted local coframing on \mathcal{X} given by the unique local section $\sigma : \mathcal{X} \rightarrow \mathcal{B}_3$, and consider the pullbacks of the structure equations (3.23) via σ . Since $\bar{\eta}^1$ is an integrable 1-form (i.e., $d\bar{\eta}^1 \equiv 0 \pmod{\bar{\eta}^1}$) but not exact, we can choose local coordinates x^1, x^2 on \mathcal{X} such that $\bar{\eta}^1 = \frac{1}{x^2} dx^1$. Then the first equation in (3.23) implies that

$$\bar{\eta}^3 \equiv \frac{1}{x^2} dx^2 \pmod{dx^1}.$$

Now, since $d\bar{\eta}^3 \not\equiv 0 \pmod{\bar{\eta}^3}$, there must exist another local coordinate x^3 , functionally independent from x^1, x^2 , such that

$$\bar{\eta}^3 = \frac{1}{x^2} \left(dx^2 - \frac{x^3}{x^2} dx^1 \right).$$

Moreover, the third equation in (3.23) implies that

$$\bar{\eta}^2 = \frac{1}{x^2} \left((dx^3 + \frac{1}{x^2} B dx^1 + \frac{1}{x^2} C \left(dx^2 - \frac{x^3}{x^2} dx^1 \right)) \right)$$

for some functions B, C on \mathcal{X} . Finally, the second equation in (3.23) implies that

$$C = \frac{1}{2} (x^2 B_{x^3} - x^3).$$

Our 3-adapted coframing now has the form

$$\begin{aligned} \bar{\eta}^1 &= \frac{1}{x^2} dx^1, \\ \bar{\eta}^2 &= \frac{1}{x^2} dx^3 + \frac{1}{(x^2)^2} B dx^1 + \frac{1}{2} \left(\frac{(x^2 B_{x^3} - x^3)}{(x^2)^2} \right) \left(dx^2 - \frac{x^3}{x^2} dx^1 \right), \\ \bar{\eta}^3 &= \frac{1}{x^2} dx^2 - \frac{x^3}{(x^2)^2} dx^1. \end{aligned} \tag{3.24}$$

The dual framing is

$$\begin{aligned} v_1 &= x^2 \frac{\partial}{\partial x^1} + x^3 \frac{\partial}{\partial x^2} - B \frac{\partial}{\partial x^3}, \\ v_2 &= x^2 \frac{\partial}{\partial x^3}, \\ v_3 &= x^2 \frac{\partial}{\partial x^2} - \frac{1}{2} (x^2 B_{x^3} - x^3) \frac{\partial}{\partial x^3}; \end{aligned}$$

setting $J = -B$ and noting that $\text{span}(v_2) = \text{span}\left(\frac{\partial}{\partial x^3}\right)$, we have:

$$\mathcal{F} = v_1 + \text{span}(v_2) = \left(x^2 \frac{\partial}{\partial x^1} + x^3 \frac{\partial}{\partial x^2} + J \frac{\partial}{\partial x^3} \right) + \text{span} \left(\frac{\partial}{\partial x^3} \right).$$

Comparing the structure equations (3.23) with those of the coframing (3.24), we see that

$$\begin{aligned} T_{12}^2 &= \frac{1}{2x^2}(x^2 J_{x^3} - 3x^3), \\ T_{13}^2 &= \frac{1}{4(x^2)^2} \left(3(x^3)^2 - 6x^2 J + 4(x^2)^2 J_{x^2} + 2x^2 x^3 J_{x^3} - 2(x^2)^3 J_{x^1 x^3} \right. \\ &\quad \left. - 2(x^2)^2 x^3 J_{x^2 x^3} + (x^2)^2 (J_{x^3})^2 - 2(x^2)^2 J J_{x^3 x^3} \right), \\ T_{23}^2 &= \frac{1}{2}(1 - x^2 J_{x^3 x^3}). \end{aligned}$$

Case 3. Suppose that $T_{23}^1 \neq 0$. (3.20) implies that, given any 2-adapted coframing $(\bar{\eta}^1, \bar{\eta}^2, \bar{\eta}^3)$, we can perform a transformation of the form (3.18) to arrive at a new 1-adapted coframing for which $T_{23}^1 = \pm 1$. Such a coframing will be called *3-adapted*. (For ease of notation, let $\varepsilon = T_{23}^1 = \pm 1$.)

Any two 3-adapted coframings on \mathcal{X} vary by a transformation of the form

$$\begin{bmatrix} \tilde{\bar{\eta}}^1 \\ \tilde{\bar{\eta}}^2 \\ \tilde{\bar{\eta}}^3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \pm 1 & 0 \\ 0 & 0 & \pm 1 \end{bmatrix}^{-1} \begin{bmatrix} \bar{\eta}^1 \\ \bar{\eta}^2 \\ \bar{\eta}^3 \end{bmatrix},$$

where the signs are chosen such that the matrix above has determinant equal to 1. The corresponding principal fiber bundle $\mathcal{B}_3 \subset \mathcal{B}_2$ has discrete fiber group $G_3 \cong \mathbb{Z}/2\mathbb{Z}$, and we can regard \mathcal{B}_3 as an (e) -structure over a 2-to-1 cover of \mathcal{X} . When restricted to \mathcal{B}_3 , the connection form β_2 becomes semi-basic; the structure equations of \mathcal{B}_3 take the form

$$\begin{aligned} d\eta^1 &= T_{13}^1 \eta^1 \wedge \eta^3 + \varepsilon \eta^2 \wedge \eta^3, \\ d\eta^2 &= T_{12}^2 \eta^1 \wedge \eta^2 + T_{13}^2 \eta^1 \wedge \eta^3 + T_{23}^2 \eta^2 \wedge \eta^3, \\ d\eta^3 &= \eta^1 \wedge \eta^2 + T_{12}^3 \eta^1 \wedge \eta^3 + T_{23}^3 \eta^2 \wedge \eta^3. \end{aligned} \tag{3.25}$$

Given any 3-adapted coframing, consider the dual framing (v_1, v_2, v_3) . The structure equations (3.25) imply that

$$[v_1, v_2] \equiv -v_3 \pmod{v_2}, \quad [v_2, v_3] \equiv -\varepsilon v_1 \pmod{v_2, v_3}, \quad [v_1, v_3] \equiv 0 \pmod{v_1, v_2, v_3}.$$

The first equation implies that $L_{\mathcal{F}^2}$ is spanned by $\{v_2, v_3\}$, and the second implies that $L_{\mathcal{F}^2}$ is not Frobenius and that \mathcal{F} is bracket-generating. Thus, this case corresponds to Case 3 of Theorem 3.4.

In order to recover the normal form of Theorem 3.4, let $(\bar{\eta}^1, \bar{\eta}^2, \bar{\eta}^3)$ be any 3-adapted local coframing on \mathcal{X} given by a local section $\sigma : \mathcal{X} \rightarrow \mathcal{B}_3$, and consider the pullbacks of the structure equations (3.25) via σ . Since $d\bar{\eta}^1 \not\equiv 0 \pmod{\bar{\eta}^1}$, Pfaff's theorem implies that we can choose local coordinates x^1, x^2, x^3 on \mathcal{X} such that

$$\bar{\eta}^1 = dx^1 - x^3 dx^2.$$

Set $T_{13}^1 = B$; this will simplify notation in what follows. The first equation in (3.25) becomes

$$(B\bar{\eta}^1 + \varepsilon\bar{\eta}^2) \wedge \bar{\eta}^3 = dx^2 \wedge dx^3,$$

which implies that

$$\begin{aligned} \bar{\eta}^2 &= \varepsilon(-B(dx^1 - x^3 dx^2) - \lambda^{-1} dx^2 + C(Hdx^2 - dx^3)), \\ \bar{\eta}^3 &= \lambda(Hdx^2 - dx^3), \end{aligned}$$

for some functions λ, C, H on \mathcal{X} with $\lambda \neq 0$. (If necessary, perform the contact transformation

$$x^1 \rightarrow x^1 - x^2 x^3, \quad x^2 \rightarrow x^3, \quad x^3 \rightarrow -x^2$$

to achieve this form.) Now the third equation in (3.25) implies that

$$-\varepsilon \lambda^2 H_{x^1} = 1.$$

Therefore, $H_{x^1} \neq 0$; $\varepsilon H_{x^1} < 0$, and

$$\lambda = \pm \frac{1}{\sqrt{-\varepsilon H_{x^1}}}.$$

The $\mathbb{Z}/2\mathbb{Z}$ freedom in the structure group G_3 may be used to eliminate the sign ambiguity in λ ; for simplicity we will take the positive square root.

Our 3-adapted coframing now has the form

$$\begin{aligned} \bar{\eta}^1 &= dx^1 - x^3 dx^2, \\ \bar{\eta}^2 &= -\varepsilon \left(\sqrt{-\varepsilon H_{x^1}} dx^2 + B(dx^1 - x^3 dx^2) + C(H dx^2 - dx^3) \right), \\ \bar{\eta}^3 &= \frac{1}{\sqrt{-\varepsilon H_{x^1}}} (H dx^2 - dx^3). \end{aligned} \tag{3.26}$$

The dual framing is

$$\begin{aligned} v_1 &= \frac{\partial}{\partial x^1} - \frac{B}{\sqrt{-\varepsilon H_{x^1}}} \left(x^3 \frac{\partial}{\partial x^1} + \frac{\partial}{\partial x^2} + H \frac{\partial}{\partial x^3} \right), \\ v_2 &= -\frac{\varepsilon}{\sqrt{-\varepsilon H_{x^1}}} \left(x^3 \frac{\partial}{\partial x^1} + \frac{\partial}{\partial x^2} + H \frac{\partial}{\partial x^3} \right), \\ v_3 &= -\sqrt{-\varepsilon H_{x^1}} \frac{\partial}{\partial x^3} + C \left(x^3 \frac{\partial}{\partial x^1} + \frac{\partial}{\partial x^2} + H \frac{\partial}{\partial x^3} \right); \end{aligned}$$

setting $J = \frac{-B}{\sqrt{-\varepsilon H_{x^1}}}$, we have:

$$\begin{aligned} \mathcal{F} &= v_1 + \text{span}(v_2) \\ &= \left(\frac{\partial}{\partial x^1} + J \left(x^3 \frac{\partial}{\partial x^1} + \frac{\partial}{\partial x^2} + H \frac{\partial}{\partial x^3} \right) \right) + \text{span} \left(x^3 \frac{\partial}{\partial x^1} + \frac{\partial}{\partial x^2} + H \frac{\partial}{\partial x^3} \right). \end{aligned}$$

Comparing the structure equations (3.25) with those of the coframing (3.26) shows that

$$\begin{aligned} C &= \frac{1}{2H_{x^1}} \sqrt{-\varepsilon H_{x^1}} \left((H_{x^1} H_{x^3} - H H_{x^1 x^3} - H_{x^1 x^2}) J \right. \\ &\quad \left. - (1 + x^3 J) H_{x^1 x^1} - (J_{x^2} + x^3 J_{x^1} + H J_{x^3}) H_{x^1} \right), \end{aligned}$$

and that

$$\begin{aligned} T_{12}^2 &= \frac{1}{2} (J_{x^2} + x^3 J_{x^1} + H J_{x^3} + J H_{x^3}), \\ T_{13}^2 &= \varepsilon C^2 H_{x^1} - H_{x^1} J_{x^3} + J^2 H_{x^1} + \frac{1}{2\sqrt{-\varepsilon H_{x^1}}} \left(2C(H_{x^1} J_{x^2} + J H_{x^1, x^2} + H_{x^1 x^1} \right. \\ &\quad \left. + 2x^3 H_{x^1} J_{x^1} + x^3 J H_{x^1 x^1} + 2H H_{x^1} J_{x^3} + H H_{x^1 x^3} J - 2H_{x^1} H_{x^3} J) \right. \\ &\quad \left. - 2x^3 C_{x^1} H_{x^1} J - 2C_{x^1} H_{x^1} - 2C_{x^3} H H_{x^1} J - 2C_{x^2} H_{x^1} J \right), \\ T_{23}^2 &= -(C_{x^2} + x^3 C_{x^1} + H C_{x^3} + C H_{x^3}) - \frac{\varepsilon}{2\sqrt{-\varepsilon H_{x^1}}} (H_{x^1 x^3} + 2J H_{x^1}), \\ T_{23}^3 &= \frac{\varepsilon}{2H_{x^1} \sqrt{-\varepsilon H_{x^1}}} (x^3 H_{x^1 x^1} - 2H_{x^1} H_{x^3} + H_{x^1 x^2} + H H_{x^1 x^3}). \end{aligned} \quad \blacksquare$$

4 Examples and questions for further study

4.1 Examples

Example 4.1 (NMR control). In quantum mechanics, a particle with spin $\frac{1}{2}$ is represented by a state vector

$$|\psi\rangle = \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \in \mathbb{C}^2$$

with $|\alpha|^2 + |\beta|^2 = 1$. In the presence of a magnetic field $\mathbf{B} = B_x \mathbf{i} + B_y \mathbf{j} + B_z \mathbf{k}$, the state vector at time t is

$$|\psi(t)\rangle = U(t)|\psi_0\rangle,$$

where $U : \mathbb{R} \rightarrow SU(2)$ is a solution of Schrödinger's equation

$$\dot{U} = -i\mathcal{H}U. \quad (4.1)$$

(See [22] for more details.) The Hamiltonian operator \mathcal{H} is given by

$$\mathcal{H} = -\gamma\hbar(B_x \cdot I_x + B_y \cdot I_y + B_z \cdot I_z),$$

where γ is the *gyromagnetic ratio* (a constant), $\hbar = \frac{h}{2\pi}$ (where h is Planck's constant), and I_x , I_y , and I_z are the *Pauli spin matrices*

$$I_x = \frac{1}{2} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad I_y = \frac{1}{2} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad I_z = \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

By multiplying the Pauli spin matrices by $-i$ as in (4.1) we obtain members of the Lie algebra of $SU(2)$. The commutation relations for $-iI_x$, $-iI_y$, and $-iI_z$ are

$$[-iI_x, -iI_y] = -iI_z, \quad [-iI_y, -iI_z] = -iI_x, \quad [-iI_z, -iI_x] = -iI_y.$$

In nuclear magnetic resonance (NMR) experiments, an ensemble of atomic nuclei of the same type can be modeled as a single spin system [14]. In such experiments, an ensemble of nuclei is subjected to a constant magnetic field \mathbf{B}_0 in the z -direction (called the *longitudinal field*), and an oscillating field \mathbf{B}_1 in the xy -plane (called the *rf field*):

$$\begin{aligned} \mathbf{B}_0 &= B_0 \mathbf{k}, \\ \mathbf{B}_1 &= B_1 (\cos(\omega t + \phi) \mathbf{i} + \sin(\omega t + \phi) \mathbf{j}), \end{aligned}$$

where ω is the frequency of the rf field and ϕ is a phase. The Hamiltonian \mathcal{H} can thus be written as

$$\mathcal{H} = -\gamma\hbar(B_0 \cdot I_z + B_1(\cos(\omega t + \phi)I_x + \sin(\omega t + \phi)I_y)).$$

In NMR experiments, the magnitude of the longitudinal field \mathbf{B}_0 is usually kept constant, while the magnitude of the rf field \mathbf{B}_1 may be used as a control variable. By transforming the coordinate system to a system that rotates at the same frequency ω as the rf field, and normalizing the constant $-\gamma\hbar B_0$ to 1, we may rewrite the Hamiltonian as

$$\mathcal{H} = I_z + u(t)I_x,$$

where $u(t) = \frac{B_1(t)}{B_0}$.

Thus Schrödinger's equation (4.1) becomes the control-affine system

$$\dot{U} = -iI_z U - (iI_x U)u \quad (4.2)$$

on the 3-dimensional state space $SU(2)$. This system corresponds to a strictly affine, rank 1 point-affine distribution

$$\mathcal{F} = v_1 + \text{span}(v_2),$$

where v_1, v_2 are the right-invariant vector fields

$$v_1 = -iI_z U, \quad v_2 = -iI_x U$$

on $SU(2)$. Let $\alpha_1, \alpha_2, \alpha_3$ denote the canonical right-invariant 1-forms on $SU(2)$ dual to the basis $-iI_x U, -iI_y U, -iI_z U$ for the right-invariant vector fields; these forms satisfy the structure equations

$$d\alpha_1 = \alpha_2 \wedge \alpha_3, \quad d\alpha_2 = \alpha_3 \wedge \alpha_1, \quad d\alpha_3 = \alpha_1 \wedge \alpha_2.$$

A 0-adapted coframing for the system (4.2) (corresponding to the choice $v_3 = -[v_1, v_2] = -iI_y U$) is given by

$$\eta^1 = \alpha_3, \quad \eta^2 = \alpha_1, \quad \eta^3 = \alpha_2.$$

This coframing has structure equations

$$d\eta^1 = \eta^2 \wedge \eta^3, \quad d\eta^2 = \eta^3 \wedge \eta^1, \quad d\eta^3 = \eta^1 \wedge \eta^2;$$

therefore this example falls into Case 3 of Theorem 3.4, with $\varepsilon = 1$. Since $T_{13}^1 = 0$, we have $J = 0$.

Let x^1, x^2, x^3 be local coordinates such that $\eta^1 = dx^1 - x^3 dx^2$. Then as in the proof of Theorem 3.4, there exist functions C, H, λ such that

$$\begin{aligned} \bar{\eta}^2 &= -\lambda^{-1} dx^2 + C(H dx^2 - dx^3), \\ \bar{\eta}^3 &= \lambda(H dx^2 - dx^3). \end{aligned}$$

(Note that $B = 0$ in this example.) The structure equations imply that

$$\begin{aligned} H &= -e^{2f(x^2, x^3)} \tan(x^1 + g(x^2, x^3)) + h(x^2, x^3), \\ C &= -e^{-f(x^2, x^3)} \sin(x^1 + g(x^2, x^3)), \\ \lambda &= e^{-f(x^2, x^3)} \cos(x^1 + g(x^2, x^3)), \end{aligned}$$

for functions f, g, h which satisfy the additional PDEs

$$\begin{aligned} e^{2f} f_{x^3} + g_{x^2} + h g_{x^3} + x^3 &= 0, \\ e^{2f} g_{x^3} + f_{x^2} + h f_{x^3} - h_{x^3} &= 0. \end{aligned}$$

Thus the invariants for this system are

$$H = -e^{2f(x^2, x^3)} \tan(x^1 + g(x^2, x^3)) + h(x^2, x^3), \quad J = 0.$$




Figure 1. Heading angle ψ in Example 4.2.

Example 4.2 (Navigation on a river). Consider a boat navigating on a river with a current; for simplicity, assume that the current runs parallel to the x -axis with constant speed c . The state of the boat is represented by its position (x, y) and heading angle ψ (see Fig. 1); thus the state space is $\mathcal{X} = \mathbb{R}^2 \times S^1$.

Due to asymmetry in the shape of the boat's hull, the current may have a rotational effect as well as a translational effect on the boat, and this effect depends on the boat's heading angle. Thus we will assume that the drift vector field has the form

$$v_1 = c \frac{\partial}{\partial x} + r(\psi) \frac{\partial}{\partial \psi}.$$

We also assume that the boat can be propelled in a forward or backward direction, corresponding to the vector field

$$v_2 = (\cos \psi) \frac{\partial}{\partial x} + (\sin \psi) \frac{\partial}{\partial y},$$

and that it can be steered to the right or left, corresponding to the vector field

$$v_3 = \frac{\partial}{\partial \psi}.$$

The navigation problem for the boat is then the control-affine system corresponding to the rank 2 point-affine distribution

$$\mathcal{F} = v_1 + \text{span}(v_2, v_3)$$

on \mathcal{X} .

Note that \mathcal{F} fails to be strictly affine precisely when $\psi = 0$ or $\psi = \pi$, since then v_1 is contained in $L_{\mathcal{F}}$. Thus our results are only applicable when the boat is *not* pointed directly upstream or downstream. Since $L_{\mathcal{F}} = \text{span}(v_2, v_3)$ is a contact distribution, this system falls into Case 3 of Corollary 1.

The dual coframing to the framing (v_1, v_2, v_3) on \mathcal{X} is

$$\bar{\eta}^1 = \frac{1}{c} (dx - (\cot \psi) dy), \quad \bar{\eta}^2 = (\csc \psi) dy, \quad \bar{\eta}^3 = d\psi - \frac{r(\psi)}{c} (dx - (\cot \psi) dy).$$

This coframing has

$$d\bar{\eta}^1 \equiv -\frac{(\csc \psi)}{c} \bar{\eta}^2 \wedge \bar{\eta}^3 \pmod{\bar{\eta}^1};$$

in order to obtain a 0-adapted coframing as in the proof of Theorem 3.3, we multiply $\bar{\eta}^2$ by $\frac{(\sin \psi)}{c}$ and $\bar{\eta}^3$ by $-(\csc^2 \psi)$, resulting in the following coframing:

$$\begin{aligned} \bar{\eta}^1 &= \frac{1}{c} (dx - (\cot \psi) dy), & \bar{\eta}^2 &= \frac{1}{c} dy, \\ \bar{\eta}^3 &= -(\csc^2 \psi) \left(d\psi - \frac{r(\psi)}{c} (dx - (\cot \psi) dy) \right). \end{aligned} \tag{4.3}$$

The local coordinate functions

$$x^1 = \frac{x}{c}, \quad x^2 = \frac{y}{c}, \quad x^3 = \cot \psi$$

are Pfaff normal form coordinates for $\bar{\eta}^1$; i.e.,

$$\bar{\eta}^1 = dx^1 - x^3 dx^2.$$

In terms of these coordinates, the coframing (4.3) becomes:

$$\bar{\eta}^1 = dx^1 - x^3 dx^2, \quad \bar{\eta}^2 = dx^2, \quad \bar{\eta}^3 = dx^3 - ((x^3)^2 + 1)R(x^3)(dx^1 - x^3 dx^2),$$

where $R(x^3) = r(\psi) = r(\cot^{-1}(x^3))$. Thus the invariants for this system are

$$J_2 = ((x^3)^2 + 1)R(x^3) = (\csc^2 \psi)r(\psi), \quad J_3 = 0.$$

4.2 Questions for further study

There are two main issues that we hope to address in future papers, both motivated by optimal control theory.

1. What metric structures (analogous to sub-Riemannian or sub-Finsler geometry for linear distributions) are appropriate for point-affine distributions, and what can we say about their geometry, geodesics, etc.? Even for linear distributions, issues such as controllability and the presence of *rigid curves* – i.e., curves with no C^1 variations whatsoever among horizontal curves – are nontrivial, and the study of optimal trajectories for sub-Riemannian and sub-Finsler metrics is quite complicated. (See, for instance, [20, 21, 9].) Agrachev and Sarychev have studied time-optimal extremals and given necessary and sufficient conditions for rigidity of trajectories for affine distributions in [2], but the consequences of imposing any type of metric structure on a point-affine distribution and the resulting geometry remain entirely unexplored. These issues are obviously important for understanding the associated control problems.
2. What can we say about the geometry of affine distributions of non-constant type, particularly for affine distributions which are not strictly affine? Normal forms for non-strictly affine distributions of rank 1 have been given by Kang and Krener [19], Kang [18], and Tall and Respondek [24], and for corank-1 affine distributions by Jakubczyk and Respondek [17], Respondek [23], and Zhitomirskii and Respondek [26]. However, intermediate ranks are important as well – for instance, for understanding second-order systems arising in mechanics, e.g.

$$\ddot{x} = \sum_{i=1}^s u^i F_i(x).$$

This system can be rewritten as the following first-order system on the cotangent bundle $T^*\mathcal{X}$, with local coordinates (x, p) :

$$\begin{aligned} \dot{x} &= p, \\ \dot{p} &= \sum_{i=1}^s u^i F_i(x). \end{aligned}$$

The associated rank s affine distribution fails to be strictly affine precisely along the submanifold $\{p = 0\}$ of stationary points. The behavior of trajectories near such points is of vital interest in control theory, so it would be useful to understand the geometry of such structures. Elkin has introduced the notion of a “ t -codistribution” in [12, 13], and we anticipate that this will be a useful tool for extending our methods to affine distributions of this type.

Acknowledgements

This work was partially supported by NSF grant DMS-0908456.

References

- [1] Agrachev A.A., Feedback-invariant optimal control theory and differential geometry. II. Jacobi curves for singular extremals, *J. Dynam. Control Systems* **4** (1998), 583–604.
- [2] Agrachev A.A., Sarychev A.V., Abnormal sub-Riemannian geodesics: Morse index and rigidity, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **13** (1996), 635–690.
- [3] Agrachev A., Zelenko I., On feedback classification of control-affine systems with one- and two-dimensional inputs, *SIAM J. Control Optim.* **46** (2007), 1431–1460, [math.OA/0502031](https://arxiv.org/abs/math.OA/0502031).
- [4] Bryant R.L., Chern S.S., Gardner R.B., Goldschmidt H.L., Griffiths P.A., Exterior differential systems, *Mathematical Sciences Research Institute Publications*, Vol. 18, Springer-Verlag, New York, 1991.
- [5] Bryant R.L., Conformal geometry and 3-plane fields on 6-manifolds, in *Developments of Cartan Geometry and Related Mathematical Problems*, *RIMS Symposium Proceedings*, Vol. 1502, Kyoto University, 2006, 1–15, [math.DG/0511110](https://arxiv.org/abs/math.DG/0511110).
- [6] Bullo F., Lewis A.D., Geometric control of mechanical systems. Modeling, analysis, and design for simple mechanical control systems, *Texts in Applied Mathematics*, Vol. 49, Springer-Verlag, New York, 2005.
- [7] Cartan É., Les systèmes de Pfaff, à cinq variables et les équations aux dérivées partielles du second ordre, *Ann. Sci. École Norm. Sup. (3)* **27** (1910), 109–192.
- [8] Clelland J.N., Moseley C.G., Sub-Finsler geometry in dimension three, *Differential Geom. Appl.* **24** (2006), 628–651, [math.DG/0406439](https://arxiv.org/abs/math.DG/0406439).
- [9] Clelland J.N., Moseley C.G., Wilkens G.R., Geometry of sub-Finsler Engel manifolds, *Asian J. Math.* **11** (2007), 699–726.
- [10] Doubrov B., Zelenko I., On local geometry of nonholonomic rank 2 distributions, [math.DG/0703662](https://arxiv.org/abs/math.DG/0703662).
- [11] Doubrov B., Zelenko I., On local geometry of rank 3 distributions with 6-dimensional square, [arXiv:0807.3267](https://arxiv.org/abs/0807.3267).
- [12] Elkin V.I., Affine control systems: their equivalence, classification, quotient systems, and subsystems. Optimization and control, 1, *J. Math. Sci. (New York)* **88** (1998), 675–721.
- [13] Elkin V.I., Reduction of nonlinear control systems. A differential geometric approach, *Mathematics and its Applications*, Vol. 472, Kluwer Academic Publishers, Dordrecht, 1999.
- [14] Ernst R., Bodenhausen G., Wokaun A., Principles of nuclear magnetic resonance in one and two dimensions, Oxford University Press, Oxford, 1987.
- [15] Gardner R.B., The method of equivalence and its applications, *CBMS-NSF Regional Conference Series in Applied Mathematics*, Vol. 58, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1989.
- [16] Ivey T.A., Landsberg J.M., Cartan for beginners: differential geometry via moving frames and exterior differential systems, *Graduate Studies in Mathematics*, Vol. 61, American Mathematical Society, Providence, RI, 2003.
- [17] Jakubczyk B., Respondek W., Feedback classification of analytic control systems in the plane, in *Analysis of Controlled Dynamical Systems (Lyon, 1990)*, *Progr. Systems Control Theory*, Vol. 8, Birkhäuser Boston, Boston, MA, 1991, 263–273.
- [18] Kang W., Extended controller form and invariants of nonlinear control systems with a single input, *J. Math. Systems Estim. Control* **6** (1996), 27–51.
- [19] Kang W., Krener A.J., Extended quadratic controller normal form and dynamic state feedback linearization of nonlinear systems, *SIAM J. Control Optim.* **30** (1992), 1319–1337.
- [20] Liu W., Sussman H.J., Shortest paths for sub-Riemannian metrics on rank-two distributions, *Mem. Amer. Math. Soc.* **118** (1995), no. 564, 104 pages.
- [21] Montgomery R., A tour of subriemannian geometries, their geodesics and applications, *Mathematical Surveys and Monographs*, Vol. 91, American Mathematical Society, Providence, RI, 2002.

-
- [22] Nielsen M., Chuang I., Quantum computation and quantum information, Cambridge University Press, Cambridge, 2000.
 - [23] Respondek W., Feedback classification of nonlinear control systems on \mathbf{R}^2 and \mathbf{R}^3 , in Geometry of Feedback and Optimal Control, *Monogr. Textbooks Pure Appl. Math.*, Vol. 207, Dekker, New York, 1998, 347–381.
 - [24] Tall I.A., Respondek W., Feedback classification of nonlinear single-input control systems with controllable linearization: normal forms, canonical forms, and invariants, *SIAM J. Control Optim.* **41** (2002), 1498–1531.
 - [25] Wilkens G.R., Centro-affine geometry in the plane and feedback invariants of two-state scalar control systems, in Differential Geometry and Control (Boulder, CO, 1997), *Proc. Sympos. Pure Math.*, Vol. 64, Amer. Math. Soc., Providence, RI, 1999, 319–333.
 - [26] Zhitomirskii M., Respondek W., Simple germs of corank one affine distributions, Singularities Symposium – Łojasiewicz 70 (Kraków, 1996; Warsaw, 1996), *Banach Center Publ.*, Vol. 44, Polish Acad. Sci., Warsaw, 1998, 269–276.