

Calvin University

Calvin Digital Commons

University Faculty Publications and Creative Works

University Faculty Scholarship

1-1-2014

An Instability Index Theory for Quadratic Pencils and Applications

Jared Bronski

University of Illinois Urbana-Champaign

Mathew A. Johnson

University of Kansas

Todd Kapitula

Calvin University

Follow this and additional works at: https://digitalcommons.calvin.edu/calvin_facultypubs



Part of the [Physics Commons](#)

Recommended Citation

Bronski, Jared; Johnson, Mathew A.; and Kapitula, Todd, "An Instability Index Theory for Quadratic Pencils and Applications" (2014). *University Faculty Publications and Creative Works*. 367.

https://digitalcommons.calvin.edu/calvin_facultypubs/367

This Article is brought to you for free and open access by the University Faculty Scholarship at Calvin Digital Commons. It has been accepted for inclusion in University Faculty Publications and Creative Works by an authorized administrator of Calvin Digital Commons. For more information, please contact digitalcommons@calvin.edu.

An Instability Index Theory for Quadratic Pencils and Applications

Jared Bronski¹, Mathew A. Johnson², Todd Kapitula³

¹ Department of Mathematics, University of Illinois at Urbana Champaign, Urbana, IL 61801, USA.

E-mail: jared@math.uiuc.edu

² Department of Mathematics, University of Kansas, Lawrence, KS 66045, USA.

E-mail: matjohn@math.ku.edu

³ Department of Mathematics and Statistics, Calvin College, Grand Rapids, MI 49546, USA.

E-mail: tmk5@calvin.edu

Received: 10 July 2012 / Accepted: 20 November 2013

Published online: 14 March 2014 – © Springer-Verlag Berlin Heidelberg 2014

Abstract: Primarily motivated by the stability analysis of nonlinear waves in second-order in time Hamiltonian systems, in this paper we develop an instability index theory for quadratic operator pencils acting on a Hilbert space. In an extension of the known theory for linear pencils, explicit connections are made between the number of eigenvalues of a given quadratic operator pencil with positive real parts to spectral information about the individual operators comprising the coefficients of the spectral parameter in the pencil. As an application, we apply the general theory developed here to yield spectral and nonlinear stability/instability results for abstract second-order in time wave equations. More specifically, we consider the problem of the existence and stability of spatially periodic waves for the “good” Boussinesq equation. In the analysis our instability index theory provides an explicit, and somewhat surprising, connection between the stability of a given periodic traveling wave solution of the “good” Boussinesq equation and the stability of the same periodic profile, but with different wavespeed, in the nonlinear dynamics of a related generalized Korteweg–de Vries equation.

Contents

1. Introduction	522
2. Preliminary Result: Spectra is Point Only	525
3. Main Result: The Instability Index Theorem	527
4. Applications to Second-Order in Time Hamiltonian Systems	532
4.1 Example: stability in (abstract) nonlinear wave equations	533
4.2 Example: periodic waves to the “good” Boussinesq equation	535
4.2.1 Case study: power law nonlinearity	545
4.2.2 Case study, continued: solitary wave and equilibrium solution limits	547
References	549

1. Introduction

When analyzing equations arising in mathematical physics and engineering, the question of stability of special families of solutions is of prominent importance as it generally determines those solutions which are most likely to be observed in physical applications. In particular, solutions which are unstable do not naturally, i.e., in absence of a controller, arise in applications except possibly as transient phenomena. Furthermore, stability analysis is often the first step in the study of finer phenomena such as transient behavior, bifurcation, and the ability to control a wave to restrict it to a stable configuration. In this paper, we are primarily motivated by recent studies into the stability of traveling wave solutions of second-order in time Hamiltonian equations of the form

$$\partial_t^2 u + \mathcal{L}_x u + \mathcal{N}(u) = 0, \quad (t, x) \in \mathbb{R}^2, \quad (1.1)$$

where \mathcal{L}_x is a self-adjoint linear operator acting on the x -variable only, and $\mathcal{N}(u)$ denotes nonlinear terms (e.g., see [2, 3, 16, 35]). A fundamental characteristic of such PDE is that they take into account weak effects of both nonlinearity and dispersion, and they arise naturally, for instance, as models for propagation of waves in nonlinear strings and in the study of bi-directional water wave propagation in the small amplitude, long wavelength regime.

In regards to the latter water wave application, an equation of particular interest in this paper (see Sect. 4.2 below) is the generalized “good” Boussinesq (gB) equation

$$\partial_t^2 u - \partial_x^2 \left(\partial_x^2 u - u + f(u) \right) = 0, \quad (1.2)$$

which is a variant of one of the equations formulated by Boussinesq in the 1870s in precisely this physical context. While the nonlinear stability and instability of solitary waves in (1.2) is by now well understood (see [4, 29]), the stability (whether linear or nonlinear) of the periodic traveling waves has received considerably less attention, and results only exist for very special classes of solutions; namely, those expressible in terms of Jacobi-elliptic functions. One of the main applications of the theory developed in this paper is an explicit connection between periodic traveling waves of (1.2) and the stability of the same traveling wave profile (but with a *different* wavespeed) in the nonlinear dynamics governed by the generalized Korteweg–de Vries (gKdV) equation

$$\partial_t u + \partial_x \left(\partial_x^2 u + f(u) \right) = 0. \quad (1.3)$$

We consider this observation as a major contribution to the theory of traveling periodic waves in (1.2). Indeed, since the stability of periodic traveling waves in gKdV equations has been under intense investigation over the last few years (see, for instance, [1, 3, 5, 6, 8, 9, 19]), this connection between the dynamics of gB and gKdV near periodic traveling waves allows one to immediately translate known results for the stability of gKdV periodic waves to results about periodic waves in gB. Applications of this connection will be illustrated in Sects. 4.2.1 and 4.2.2 below.

Suppose that $u(x - ct)$ is a traveling wave solution to (1.1). When considering small perturbations to this wave of the form $e^{\lambda t} v(x - ct)$, $\lambda \in \mathbb{C}$, we are naturally led to quadratic spectral problems of the form

$$\lambda^2 v - 2c\lambda \partial_x v + \left(c^2 \partial_x^2 + \mathcal{L}_x + \mathcal{N}'(u) \right) v = 0.$$

More generally, when considering the spectral stability of the given nonlinear dispersive wave u , it becomes important to understand the spectrum of quadratic operator pencils of the form

$$\mathcal{P}_2(\lambda) := \mathcal{A} + \lambda\mathcal{B} + \lambda^2\mathcal{C}, \quad (1.4)$$

where the operators \mathcal{A} and \mathcal{C} are self-adjoint on some Hilbert space X , endowed with an inner-product $\langle \cdot, \cdot \rangle$, and \mathcal{B} is skew-symmetric on X . The class of perturbations considered in our applications naturally leads us to assume that the domain of each of the operators $\mathcal{A}, \mathcal{B}, \mathcal{C}$ is dense in X and that they each enjoy a particular compactness property which guarantees that the spectrum of $\mathcal{P}_2(\lambda)$, i.e., the collection of values λ for which $\mathcal{P}_2(\lambda)$ fails to be boundedly invertible, is composed of point spectrum only, that each eigenvalue has finite algebraic multiplicity, and the only accumulation point of the eigenvalues is infinity (see Lemma 2.1 below for a precise statement). In this paper $\sigma(\mathcal{P}_2)$ will denote the collection of all eigenvalues for the pencil.

Due to its clear connection and importance in analyzing the spectral stability of traveling wave solutions of equations of the form (1.1), our main theoretical results concern extending many previously known results regarding the number of eigenvalues of \mathcal{P}_2 with positive real part. For the reader's convenience, we now briefly recall the relevant known results. The spectrum was studied in [33] under the assumption that \mathcal{A} has compact resolvent, and \mathcal{B}, \mathcal{C} are bounded and positive semi-definite. The operators \mathcal{B}, \mathcal{C} are not assumed to have any symmetry properties, however. While it is not shown in that paper, by Lemma 2.1 it is then known that the pencil only possesses point eigenvalues, each of which has finite algebraic multiplicity. It is shown in that paper that if \mathcal{A} is positive semi-definite, then all of the spectrum is located in the closed left-half of the complex plane. If \mathcal{A} is not definite, it is shown that if $\mathcal{C} = \mathcal{I}$ and \mathcal{B} is positive definite, then the total number of eigenvalues in the closed right-half of the complex-plane is equal to the number of negative eigenvalues of \mathcal{A} . As is seen in, e.g., Sect. 3, it is not necessary that $\mathcal{C} = \mathcal{I}$ in order for that result to hold; indeed, all that is needed is that \mathcal{C} be positive-definite. In [30] similar results are shown regarding the number of eigenvalues with positive real part under the assumption that $\mathcal{C} = \mathcal{I}$ and \mathcal{B} is positive semi-definite. The spectrum of operators of the form \mathcal{P}_2 was studied in [15] under the assumptions that both \mathcal{A}, \mathcal{C} are positive definite and self-adjoint, while \mathcal{B} is a negative definite self-adjoint operator (also see [27] for a generalization). Under assumptions different than those given in Lemma 2.1 (in particular, both \mathcal{A} and \mathcal{C} are assumed to be compact) it is shown therein that in $\sigma(\mathcal{P}_2)$ there is an infinite number of positive eigenvalues which have zero as a limit point. Finally, the matrix-valued version of the pencil was studied in [7, 27] under the assumption that $\mathcal{C} = \mathcal{I}$ and \mathcal{A}, \mathcal{B} self-adjoint. Therein a parity index is given relating the number of eigenvalues with positive real part to the number of negative eigenvalues of \mathcal{A} . This result strongly depends upon the fact that the operators are matrix-valued, and consequently the Hilbert space under question is finite-dimensional.

One of the main goals of this paper is thus to extend the previous theory regarding the number of eigenvalues (counting multiplicity) with positive real part; see Sect. 3 below. Throughout this analysis, two underlying assumptions will be:

- (a) $n(\mathcal{A}), n(\mathcal{C}) < +\infty$, where $n(\mathcal{S})$ refers to the number of negative eigenvalues (counting multiplicity) of the self-adjoint operator \mathcal{S}
- (b) \mathcal{C} is invertible, i.e., $z(\mathcal{C}) = 0$, where $z(\mathcal{S}) = \dim[\ker(\mathcal{S})]$ for the self-adjoint operator \mathcal{S} .

Under these assumptions we have the intuition that the sum $n(\mathcal{A}) + n(\mathcal{C})$ acts as an upper bound on the total number of eigenvalues of \mathcal{P}_2 with positive real part. Indeed, if the linear term in the pencil is dropped, then the quadratic pencil becomes

$$\mathcal{P}_2(\lambda) = \mathcal{A} + \lambda^2 \mathcal{C}.$$

The spectral theory of this quadratic pencil was studied by Grillakis [12]. By considering the Krein signature of the purely imaginary eigenvalues, and using the Euler characteristic of the projective sphere, he was able to show that the intuitive upper bound is actually the true upper bound. Another proof follows by linearizing the pencil. When the linear term is dropped the quadratic pencil is equivalent to the linear pencil

$$\left[\begin{pmatrix} \mathcal{A} & 0 \\ 0 & \mathcal{C}^{-1} \end{pmatrix} - \lambda \begin{pmatrix} 0 & \mathcal{I} \\ -\mathcal{I} & 0 \end{pmatrix} \right] \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Note that the first term in the linear pencil is self-adjoint, whereas the second term is skew-symmetric. Linear pencils of this form have been well-studied (e.g., see [18, 25, 26, 32] and the references therein), and for this problem it has firmly been established that the intuition is indeed correct. The technical difficulty to establish the result for the full pencil is then the inclusion of the linear term, and it is overcome in the establishment of Lemma 3.1 below. In particular, in the case where $\mathcal{P}_2(\lambda)$ arises in the stability analysis of a given periodic traveling wave (as described previously), a sufficient condition for spectral stability is that $n(\mathcal{A}) + n(\mathcal{C}) = 0$.

In order to achieve an equality relating the number of eigenvalues of \mathcal{P}_2 with positive real part to spectral properties of the operators \mathcal{A} , \mathcal{B} , and \mathcal{C} themselves, two additional factors must be taken into account:

- (a) the effect of \mathcal{A} having a nontrivial kernel
- (b) purely imaginary eigenvalues having a negative Krein index (see Sect. 3).

The first factor arises because in applications the presence of symmetries yields the existence of a nontrivial kernel, while the consideration of the second factor is necessary in order to remove the intuitive inequality and make it an equality.

The paper is organized as follows. In Sect. 2 it is shown that under general assumptions, which are natural for the applications we have in mind, there is no essential spectrum for the quadratic operator pencil \mathcal{P}_2 ; in other words, there will be only point eigenvalues, and each eigenvalue will have finite algebraic multiplicity. The proof of this result easily generalizes to polynomial operators of arbitrary order; see Remark 2.2 below. In Sect. 3 the main theoretical result of the paper relating the number of eigenvalues of \mathcal{P}_2 with positive real part to the spectral properties of \mathcal{A} , \mathcal{B} , and \mathcal{C} , are stated and proved. Finally, in Sect. 4 we apply the theoretical results developed in Sects. 2 and 3 to the study of the spectral and orbital stability of nonlinear waves to second-order in time Hamiltonian systems. The first application develops a general theoretical result concerning the stability of steady states in abstract nonlinear wave equations; see Sect. 4.1. The second application examines the spectral and nonlinear (orbital) stability of periodic traveling waves in the “good” Boussinesq equation (1.2); see Sect. 4.2. As described above, of particular interest in our study of the Boussinesq equation is that we establish in Lemma 4.2 and Theorem 4.8 a rigorous connection between a given stationary periodic wave u of the equation

$$\partial_t^2 u - 2c \partial_{tX}^2 u + \partial_X^2 \left(\partial_X^2 u - (1 - c^2)u + f(u) \right) = 0, \quad |c| < 1,$$

corresponding to the traveling wave $u(x - ct)$ in the gB, and the stability of the same stationary wave profile in the gKdV,

$$\partial_t u + \partial_x \left(\partial_x^2 - \sqrt{1 - c^2} u + f(u) \right) = 0.$$

To illustrate the power of this connection, in Sect. 4.2.1 we translate recent results by Bronski et al. [6] concerning the stability of periodic KdV waves with power-law nonlinearity to results for the corresponding Boussinesq equation, allowing one to see the complete stability picture of periodic waves in gB in the cases considered. In Sect. 4.2.2, using known asymptotic analyses worked out in the context of stability theory for gKdV waves (see [5]), we analyze the stability of periodic waves with power-law nonlinearity which are near (in an appropriate sense) either the solitary wave or equilibrium solutions of gB. The analysis near the solitary wave makes a beautiful connection with the known nonlinear stability/instability results for the nearby solitary waves. Similarly, the analysis near the equilibrium solution provides new insights into the transitions to instability in the periodic context; in particular, we find that for a given power-law nonlinearity stable periodic traveling wave solutions always exist, even when no stable solitary wave exists.

2. Preliminary Result: Spectra is Point Only

Notation: In this paper, and particularly in this and the subsequent section, the notion of *matrix representation* of a self-adjoint operator S constrained to a subspace E , i.e., $S|_E$, will often be used. Let $\{e_1, \dots, e_n\}$ be a basis for E , and let $P_E : X \mapsto E$ be the orthogonal projection. If the basis is orthonormal, then one can write

$$P_E = \sum_{j=1}^n \langle \cdot, e_j \rangle e_j.$$

Setting $S|_E = P_E S P_E : E \mapsto E$, the quadratic form

$$\langle u, S|_E u \rangle = \langle P_E u, P_E S P_E u \rangle = \mathbf{c} \cdot \mathbf{S} \mathbf{c},$$

where $P_E u = \sum c_j e_j$, $\mathbf{c} = (c_1, \dots, c_n)^T$, and the symmetric matrix $\mathbf{S} \in \mathbb{R}^{n \times n}$ is given by $S_{ij} = \langle e_i, S e_j \rangle$. The matrix \mathbf{S} is precisely the matrix representation for the self-adjoint operator $S|_E$. In the rest of this paper the symbol $S|_E$ will be used to represent the matrix representation \mathbf{S} of S constrained to operate on the subspace E .

The goal of this section is to demonstrate that for the quadratic pencil of (1.4), i.e.,

$$\mathcal{P}_2(\lambda) = \mathcal{A} + \lambda \mathcal{B} + \lambda^2 \mathcal{C},$$

there is point spectra only, that each eigenvalue has finite multiplicity, and infinity is the only possible limit point of the eigenvalues. The result requires the use of [31, Theorem 12.9], in which it is shown that the spectrum has the desired properties for the polynomial operator

$$\mathcal{P}_n(\lambda) = \mathcal{I} + \sum_{j=1}^n \lambda^j \mathcal{A}_j$$

if each operator \mathcal{A}_j , $j = 1, \dots, n$, is compact. With this result in mind, first assume that \mathcal{A} is invertible, and rewrite the eigenvalue problem as

$$\mathcal{A}(\mathcal{I} + \lambda\mathcal{A}^{-1}\mathcal{B} + \lambda^2\mathcal{A}^{-1}\mathcal{C})u = 0.$$

Of course, since \mathcal{A} is nonsingular this problem is equivalent to

$$(\mathcal{I} + \lambda\mathcal{A}^{-1}\mathcal{B} + \lambda^2\mathcal{A}^{-1}\mathcal{C})u = 0.$$

If the operators $\mathcal{A}^{-1}\mathcal{B}$ and $\mathcal{A}^{-1}\mathcal{C}$ are both compact, then the result immediately follows from [31, Theorem 12.9].

On the other hand, now suppose that $\ker(\mathcal{A})$ is nontrivial, but that \mathcal{A} has compact resolvent. The proof of the desired result will now be accomplished via the construction and evaluation of a generalized *Krein matrix*, which was recently introduced in a general form in [22]. Let $P_{\mathcal{A}} : X \mapsto \ker(\mathcal{A})$ be the orthogonal projection. Writing $u = a + a^\perp$, where $a \in \ker(\mathcal{A})$ and $a^\perp \in \ker(\mathcal{A})^\perp$, the eigenvalue problem becomes

$$\mathcal{P}_2(\lambda)a + \mathcal{P}_2(\lambda)a^\perp = 0. \tag{2.1}$$

Defining the complementary projection $P_{\mathcal{A}}^\perp := \mathcal{I} - P_{\mathcal{A}}$, applying this projection to (2.1), and solving for $a^\perp = P_{\mathcal{A}}^\perp a^\perp$ yields

$$a^\perp = -(P_{\mathcal{A}}^\perp \mathcal{P}_2(\lambda) P_{\mathcal{A}}^\perp)^{-1} P_{\mathcal{A}}^\perp \mathcal{P}_2(\lambda) a. \tag{2.2}$$

In the formulation of (2.2) it is implicitly being assumed that $\mathcal{P}_2(\lambda)a^\perp \neq 0$. If $\mathcal{P}_2(\lambda)a^\perp = 0$, then λ is an eigenvalue whose eigenfunction is in $\ker(\mathcal{A})^\perp$: this follows immediately from (2.1) upon setting $a = 0$. Since $\mathcal{P}_2(\lambda)a = 0$, the potential pole singularity for such a λ is removable. If the inner-product with a is now taken in (2.1), then it is seen that

$$\langle a, \mathcal{P}_2(\lambda)a \rangle + \langle a^\perp, \mathcal{P}_2(\lambda)a \rangle = 0, \tag{2.3}$$

where

$$\mathcal{P}_2(\lambda)^a = \mathcal{A} - \bar{\lambda}\mathcal{B} + \bar{\lambda}^2\mathcal{C}$$

is the adjoint operator for the original pencil. Note that the fact that \mathcal{A}, \mathcal{C} are self-adjoint and \mathcal{B} is skew-symmetric was used in this formulation of the adjoint pencil. Substituting the expression for s^\perp given in (2.2) into (2.3) yields the linear system

$$\mathbf{K}_2(\lambda)\mathbf{x} = \mathbf{0}, \quad \mathbf{K}_2(\lambda) := \mathcal{P}_2(\lambda)|_{\ker(\mathcal{A})} - (P_{\mathcal{A}}^\perp \mathcal{P}_2(\lambda) P_{\mathcal{A}}^\perp)^{-1}|_{P_{\mathcal{A}}^\perp \mathcal{P}_2(\lambda)[\ker(\mathcal{A})]}. \tag{2.4}$$

Here

$$\mathbf{x} \in \mathbb{C}^{z(\mathcal{A})}, \quad \mathbf{K}_2(\lambda) \in \mathbb{C}^{z(\mathcal{A}) \times z(\mathcal{A})},$$

and

$$\mathcal{P}_2(\lambda)[\ker(\mathcal{A})] := \{\mathcal{P}_2(\lambda)a : a \in \ker(\mathcal{A})\}.$$

The matrix $\mathbf{K}_2(\lambda)$ is known as the Krein matrix.

Eigenvalues for the pencil are found either via $\mathbf{x} = \mathbf{0}$, which means that if λ is an eigenvalue, then the associated eigenfunction satisfies $u \in \ker(\mathcal{A})^\perp$, or $\det[\mathbf{K}_2(\lambda)] = 0$.

The quest for eigenvalues is now a search for the zeros of the determinant of the Krein matrix. The poles of the Krein matrix are the eigenvalues of the operator

$$P_{\mathcal{A}}^{\perp} \mathcal{P}_2(\lambda) P_{\mathcal{A}}^{\perp} = P_{\mathcal{A}}^{\perp} \mathcal{A} P_{\mathcal{A}}^{\perp} + \lambda P_{\mathcal{A}}^{\perp} \mathcal{B} P_{\mathcal{A}}^{\perp} + \lambda^2 P_{\mathcal{A}}^{\perp} \mathcal{C} P_{\mathcal{A}}^{\perp} : \ker(\mathcal{A})^{\perp} \mapsto \ker(\mathcal{A})^{\perp}.$$

Since $P_{\mathcal{A}}^{\perp} \mathcal{A} P_{\mathcal{A}}^{\perp} : \ker(\mathcal{A})^{\perp} \mapsto \ker(\mathcal{A})^{\perp}$ is invertible, from the earlier argument in this section we know that the operator $P_{\mathcal{A}}^{\perp} \mathcal{P}_2(\lambda) P_{\mathcal{A}}^{\perp}$ has point spectra only, each eigenvalue has finite multiplicity, and infinity is the only possible limit point of the eigenvalues. Thus, one can say that $\det[\mathbf{K}_2(\lambda)] : \mathbb{C} \mapsto \mathbb{C}$ is meromorphic, each singularity is a pole of finite order, and the only possible accumulation point of the poles is infinity. Since $\det[\mathbf{K}_2(\lambda)]$ is meromorphic, it is then known that each of its zeros is of finite order, and that the only possible accumulation point of the zeros is infinity. Finally, it was demonstrated in [22] that for linear pencils the order of the zero of $\det[\mathbf{K}_2(\lambda)]$ is equal to the algebraic multiplicity of the eigenvalue. The proof of that result easily carries over to quadratic pencils, and will be left for the interested reader.

Lemma 2.1. *Let $P_{\mathcal{A}} : X \mapsto \ker(\mathcal{A})$ be the orthogonal projection, and let $P_{\mathcal{A}}^{\perp} = \mathcal{I} - P_{\mathcal{A}}$ be the complementary projection. Suppose that the operators*

$$(P_{\mathcal{A}}^{\perp} \mathcal{A} P_{\mathcal{A}}^{\perp})^{-1} P_{\mathcal{A}}^{\perp} \mathcal{B} P_{\mathcal{A}}^{\perp}, (P_{\mathcal{A}}^{\perp} \mathcal{A} P_{\mathcal{A}}^{\perp})^{-1} P_{\mathcal{A}}^{\perp} \mathcal{C} P_{\mathcal{A}}^{\perp} : \ker(\mathcal{A})^{\perp} \mapsto \ker(\mathcal{A})^{\perp}$$

are both compact. Then the spectrum of the quadratic pencil $\mathcal{A} + \lambda \mathcal{B} + \lambda^2 \mathcal{C}$ is point spectra only. Furthermore, each eigenvalue has finite multiplicity, and infinity is the only possible limit point of the eigenvalues.

Remark 2.2. Note that the proof of Lemma 2.1 did not require that any of the operators have a symmetry property. Thus, this lemma can be thought of as a general result about quadratic pencils. Indeed, although it will not be proven here, the above argument can be extended to show that for n th-order polynomial pencils of the form

$$\mathcal{P}_n(\lambda) = \sum_{j=0}^n \lambda^j \mathcal{A}_j,$$

if each of the operators $(P_{\mathcal{A}_0}^{\perp} \mathcal{A}_0 P_{\mathcal{A}_0}^{\perp})^{-1} P_{\mathcal{A}_0}^{\perp} \mathcal{A}_j P_{\mathcal{A}_0}^{\perp}$ is compact for $j = 1, \dots, n$, then the spectrum for this pencil will have exactly the same properties as that for the quadratic pencil.

3. Main Result: The Instability Index Theorem

The goal here is to derive an instability index theorem for the quadratic pencil. In applications it may be the case that \mathcal{C}^{-1} is bounded and not compact, and/or \mathcal{B} is not bounded. In order to overcome these technical difficulties (which are associated with the proof only), take a positive definite self-adjoint operator \mathcal{S} which has a compact inverse, and consider the quadratic pencil

$$\widehat{\mathcal{P}}_2(\lambda) := \widehat{\mathcal{A}} + \lambda \widehat{\mathcal{B}} + \lambda^2 \widehat{\mathcal{C}}.$$

The operators in this new pencil are related to those of the original pencil by

$$\widehat{\mathcal{A}} := \mathcal{S}^{-1} \mathcal{A} \mathcal{S}^{-1}, \quad \widehat{\mathcal{B}} := \mathcal{S}^{-1} \mathcal{B} \mathcal{S}^{-1}, \quad \widehat{\mathcal{C}} := \mathcal{S}^{-1} \mathcal{C} \mathcal{S}^{-1},$$

so that

$$\mathcal{P}_2(\lambda) = \mathcal{S}\widehat{\mathcal{P}}_2(\lambda)\mathcal{S}.$$

The working assumption is that \mathcal{B} , \mathcal{C} are \mathcal{S} -compact, while \mathcal{A} has \mathcal{S} -compact resolvent, where an operator \mathcal{T} is said to be \mathcal{S} -compact if the operator $\mathcal{S}^{-1}\mathcal{T}\mathcal{S}^{-1}$ is compact, and is said to have \mathcal{S} -compact resolvent if the inverse $\mathcal{S}\mathcal{T}^{-1}\mathcal{S}$ (defined to act on the range space) is compact.

Define the space $X_{\mathcal{S}}$ by

$$X_{\mathcal{S}} := \{u \in X : \langle \mathcal{S}u, \mathcal{S}u \rangle < \infty\}.$$

It will be assumed that $X_{\mathcal{S}} \subset X$ is dense. Since \mathcal{S}^{-1} is compact, and therefore bounded, it is clear that $\sigma(\widehat{\mathcal{P}}_2) \subset \sigma(\mathcal{P}_2)$ on X . It is also clear that $\sigma(\mathcal{P}_2)$ when considered on the space $X_{\mathcal{S}}$ is a subset of $\sigma(\widehat{\mathcal{P}}_2)$ when considered on the space X . In other words, it is true that

$$\sigma(\mathcal{P}_2) \text{ on } X_{\mathcal{S}} \subset \sigma(\widehat{\mathcal{P}}_2) \text{ on } X \subset \sigma(\mathcal{P}_2) \text{ on } X.$$

Since $X_{\mathcal{S}} \subset X$ is dense, by [34, Proposition 3.2] it will be the case that

$$\sigma(\mathcal{P}_2) \text{ on } X_{\mathcal{S}} = \sigma(\widehat{\mathcal{P}}_2) \text{ on } X.$$

Finally, since \mathcal{S}^{-1} is bounded, it is true that $n(\mathcal{A}) = n(\widehat{\mathcal{A}})$ and $n(\mathcal{C}) = n(\widehat{\mathcal{C}})$. In conclusion, from this point forward the ‘‘hat’’s associated with the operators can be dropped, and it will be assumed that the operators satisfy \mathcal{A}^{-1} , \mathcal{C} , \mathcal{B} are compact.

The instability index theorem for the quadratic pencil will be proven by constructing an equivalent linear pencil, and then deriving an index theorem for that linear pencil. Upon setting $w = (u, \lambda\mathcal{C}u)^T$, the quadratic pencil (1.4) is linearized to become

$$(\mathcal{L} - \lambda\mathcal{J}^{-1})w = 0, \tag{3.1}$$

where

$$\mathcal{L} = \begin{pmatrix} \mathcal{A} & 0 \\ 0 & \mathcal{C}^{-1} \end{pmatrix}, \quad \mathcal{J} = \begin{pmatrix} 0 & \mathcal{I} \\ -\mathcal{I} & -\mathcal{B} \end{pmatrix}.$$

Here is where the assumption that \mathcal{C} be invertible comes into play. Since \mathcal{B} is skew-symmetric, (3.1), and consequently the pencil (1.4), is formally equivalent to the Hamiltonian eigenvalue problem

$$\mathcal{J}\mathcal{L}w = \lambda w, \tag{3.2}$$

where \mathcal{J} is skew-symmetric and \mathcal{L} is self-adjoint. Note that in this formulation

- (a) \mathcal{L} has compact resolvent
- (b) \mathcal{J} has bounded inverse
- (c) the spectrum satisfies the symmetry $\{\lambda, -\bar{\lambda}\} \subset \sigma(\mathcal{J}\mathcal{L})$, and if all of the operators have zero imaginary part, the symmetry becomes $\{\pm\lambda, \pm\bar{\lambda}\} \subset \sigma(\mathcal{J}\mathcal{L})$.

Since \mathcal{C} is invertible, it is clear that for nonzero λ the two eigenvalue problems have identical eigenvalues; furthermore, the geometric multiplicities match. As it will be seen, it is also the case that the algebraic multiplicities of these eigenvalues also coincide. We will show that this is true at the origin: the proof will clearly generalize to the case of a nonzero eigenvalue. We must consider the structure of $\text{gker}(\mathcal{J}\mathcal{L})$ and the manner in which it relates to $\text{gker}(\mathcal{P}_2(0))$.

First consider $\text{gker}(\mathcal{P}_2(0))$. It is clear that $\ker(\mathcal{P}_2(0)) = \ker(\mathcal{A})$. Following Markus [31], generalized eigenfunctions are found by solving

$$\mathcal{P}_2(0)a_1 + \mathcal{P}'_2(0)a_0 = 0, \quad a_0 \in \ker(\mathcal{A}).$$

This is equivalent to solving

$$\mathcal{A}a_1 = -\mathcal{B}a_0, \tag{3.3}$$

which by the Fredholm alternative has a nontrivial solution if and only if $\mathcal{B}a_0 \in \ker(\mathcal{A})^\perp$. Since \mathcal{B} is skew-symmetric, it is not unreasonable to assume that $\mathcal{B}|_{\ker(\mathcal{A})} = \mathbf{0}$. In this case there is a solution to (3.3) for any $a_0 \in \ker(\mathcal{A})$, and a full set of associated eigenfunctions is given by $\mathcal{A}^{-1}\mathcal{B}\ker(\mathcal{A})$, where the obvious notation

$$\mathcal{A}^{-1}\mathcal{B}\ker(\mathcal{A}) := \{\mathcal{A}^{-1}\mathcal{B}a : a \in \ker(\mathcal{A})\}$$

is being used. The next set of eigenfunctions is found by solving

$$\mathcal{P}_2(0)a_2 + \mathcal{P}'_2(0)a_1 + \frac{1}{2}\mathcal{P}''_2(0)a_0 = 0, \quad a_0 \in \ker(\mathcal{A}), \quad a_1 = -\mathcal{A}^{-1}\mathcal{B}a_0 \in \mathcal{A}^{-1}\mathcal{B}\ker(\mathcal{A}).$$

This equation is equivalent to

$$\mathcal{A}a_2 = -\mathcal{B}a_1 - \mathcal{C}a_0 = -(\mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B})a_0. \tag{3.4}$$

Again using the Fredholm alternative, it is seen that there is a nontrivial solution to (3.4) if and only if $(\mathcal{B}\mathcal{A}^{-1}\mathcal{B} - \mathcal{C})a_0 \in \ker(\mathcal{A})^\perp$. Upon using the fact that for any $a \in \ker(\mathcal{A})$,

$$\langle \mathcal{B}(\mathcal{A}^{-1}\mathcal{B}a_0), a \rangle = -\langle \mathcal{A}^{-1}\mathcal{B}a_0, \mathcal{B}a \rangle,$$

it is seen that if

$$\mathbf{D} := (\mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B})|_{\ker(\mathcal{A})}$$

is nonsingular, then $(\mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B})a_0 \notin \ker(\mathcal{A})^\perp$ for any $a_0 \in \ker(\mathcal{A})$. It will henceforth be assumed that \mathbf{D} is nonsingular.

Now consider $\text{gker}(\mathcal{J}\mathcal{L})$. It is clear that $\ker(\mathcal{L}) = (\ker(\mathcal{A}), 0)^\text{T}$. Since

$$\mathcal{J}^{-1}\ker(\mathcal{L}) = \left\{ \begin{pmatrix} -\mathcal{B}a_0 \\ a_0 \end{pmatrix} : a_0 \in \ker(\mathcal{A}) \right\}, \tag{3.5}$$

upon writing $w = (u, v)^\text{T}$ the generalized eigenfunctions are found by solving

$$\mathcal{A}u = -\mathcal{B}a_0, \quad \mathcal{C}^{-1}v = a_0.$$

The first equation is precisely (3.3), which was seen to have a solution for any $a_0 \in \ker(\mathcal{A})$. Consequently, there is a generalized eigenspace at $\lambda = 0$ which is given by $\{(-\mathcal{A}^{-1}\mathcal{B}a_0, \mathcal{C}a_0)^T : a_0 \in \ker(\mathcal{A})\}$. Since

$$\mathcal{J}^{-1} \begin{pmatrix} -\mathcal{A}^{-1}\mathcal{B}a_0 \\ \mathcal{C}a_0 \end{pmatrix} = - \begin{pmatrix} (\mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B})a_0 \\ \mathcal{A}^{-1}\mathcal{B}a_0 \end{pmatrix},$$

the next set of generalized eigenfunctions is found by solving

$$\mathcal{A}u = -(\mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B})a_0, \quad \mathcal{C}^{-1}v = -\mathcal{A}^{-1}\mathcal{B}a_0.$$

The first equation is precisely (3.4), which was seen to have no solution under the assumption that \mathbf{D} is nonsingular.

It is now seen that regarding the algebraic multiplicity of the eigenvalue the Hamiltonian linearization (3.2) is equivalent to the pencil at $\lambda = 0$. Following the same argument for nonzero λ it is not difficult to check that this equivalence continues to hold; namely, the location of the eigenvalues, and their multiplicities, are the same for the two systems.

We are now ready to derive the instability index for the linear Hamiltonian system (3.2). We must first construct the appropriate closed subspace on which both \mathcal{J} , \mathcal{L} are nonsingular. Recalling that $P_{\mathcal{A}}^{\perp} : X \mapsto \ker(\mathcal{A})^{\perp}$ is the orthogonal projection, let $\Pi_{\mathcal{L}}^{\perp} : X \times X \mapsto \ker(\mathcal{A})^{\perp} \times X = \ker(\mathcal{L})^{\perp}$ be given by

$$\Pi_{\mathcal{L}}^{\perp} := \begin{pmatrix} P_{\mathcal{A}}^{\perp} & 0 \\ 0 & \mathcal{I} \end{pmatrix};$$

in other words, for $w = (u, v) \in X \times X$ it is true that $\Pi_{\mathcal{L}}^{\perp}w = (P_{\mathcal{A}}^{\perp}u, v)$. Define another orthogonal projection by

$$\Pi_{\mathcal{J}^{-1} \ker(\mathcal{L})}^{\perp} : X \times X \mapsto [\mathcal{J}^{-1} \ker(\mathcal{L})]^{\perp}. \tag{3.6}$$

Because $\ker(\mathcal{L}) \perp \mathcal{J}^{-1} \ker(\mathcal{L})$, the projections $\Pi_{\mathcal{L}}^{\perp}$, $\Pi_{\mathcal{J}^{-1} \ker(\mathcal{L})}^{\perp}$ commute. Upon setting $\Pi := \Pi_{\mathcal{L}}^{\perp} \Pi_{\mathcal{J}^{-1} \ker(\mathcal{L})}^{\perp}$ (note that Π being the composition of self-adjoint commuting operators implies that it too is self-adjoint), nonzero eigenvalues for the linearization (3.2) are found by solving

$$\Pi \mathcal{J} \Pi \cdot \Pi \mathcal{L} \Pi \cdot \Pi w = \lambda \Pi w, \quad \Pi w \in [\ker(\mathcal{L}) \oplus \mathcal{J}^{-1} \ker(\mathcal{L})]^{\perp} \tag{3.7}$$

(e.g., see [8, Section 2]). This is the eigenvalue problem to be studied in the rest of this section.

The goal is to now count the total number of eigenvalues in the open right-half of the complex plane (counting multiplicity), along with those eigenvalues on the imaginary axis which have negative Krein index. For each $\lambda \in i\mathbb{R}$ let E_{λ} denote the generalized eigenspace. The negative Krein index of the eigenvalue for the linearized system (3.2) is given by

$$k_i^-(\lambda) := n(\mathcal{L}|_{E_{\lambda}}),$$

and if $k_i^-(\lambda) = 0$, then the eigenvalue is (often) said to have positive Krein signature. Let us now relate this definition to a definition using the quadratic pencil. By the definition of w leading to the linearization (3.1) one has that

$$\mathcal{L}|_{E_{\lambda}} = (\mathcal{A} + |\lambda|^2\mathcal{C})|_{\Pi_1 E_{\lambda}} = (\mathcal{A} - \lambda^2\mathcal{C})|_{\Pi_1 E_{\lambda}},$$

where $\Pi_1 : X \times X \mapsto X$ is the projection onto the first component, i.e., $\Pi_1(u, v)^T = u$, and the second equality follows from the fact that $\lambda \in i\mathbb{R}$. Now, λ being an eigenvalue with associated eigenfunction u means that

$$\mathcal{P}_2(\lambda)u = 0 \Rightarrow \mathcal{A}u = -\lambda\mathcal{B}u - \lambda^2\mathcal{C}u,$$

which in turn implies

$$(\mathcal{A} - \lambda^2\mathcal{C})|_{\Pi_1 E_\lambda} = -\lambda\mathcal{P}'_2(\lambda)|_{\Pi_1 E_\lambda}.$$

In conclusion, the negative Krein index for the quadratic pencil is defined to be

$$k_i^-(\lambda) := n(-\lambda\mathcal{P}'_2(\lambda)|_{E_\lambda}), \tag{3.8}$$

where now $E_\lambda = \text{gker}(\mathcal{P}_2(\lambda))$, i.e., the generalized eigenspace of the quadratic pencil $\mathcal{P}_2(\lambda)$ associated with the eigenvalue $\lambda \in i\mathbb{R}$.

We are now ready to derive the index formula. For the eigenvalue problem (3.7) let k_r represent the number of positive real-valued eigenvalues (counting multiplicity), and let k_c be the number of complex-valued eigenvalues (counting multiplicity) with positive real part. Furthermore, let the total negative Krein index be given by

$$k_i^- := \sum_{\sigma(\mathcal{P}_2(\lambda)) \cap i\mathbb{R}} k_i^-(\lambda).$$

Regarding the eigenvalue problem (3.7) it is known that $(\Pi\mathcal{L}\Pi)^{-1}$ is compact, and that the operator $\Pi\mathcal{J}\Pi$ is bounded with bounded inverse. Indeed, since one can write

$$\Pi\mathcal{J}\Pi = \Pi \begin{pmatrix} 0 & \mathcal{I} \\ -\mathcal{I} & 0 \end{pmatrix} \Pi + \Pi \begin{pmatrix} 0 & 0 \\ 0 & -\mathcal{B} \end{pmatrix} \Pi,$$

one actually has that both $\Pi\mathcal{J}\Pi$ and $(\Pi\mathcal{J}\Pi)^{-1}$ can be written as the sum of a bounded operator and a compact operator. Thus, upon using the fact that compact operators are uniformly approximated by matrices, when computing an index which takes into the account the (finite) number of negative directions of an operator it is sufficient to consider the case of matrices only. For the eigenvalue problem (3.7) when the operators are matrices it is known from [18, 23] that

$$k_r + k_c + k_i^- = n(\Pi\mathcal{L}\Pi). \tag{3.9}$$

Since all of the quantities are integer-valued, by taking the limit one deduces that the result holds for the full operators.

Before stating the final result, the quantity $n(\Pi\mathcal{L}\Pi)$ must be computed in terms of the original operator \mathcal{L} . It is known (e.g., see [24, Index Theorem]) that

$$n(\Pi\mathcal{L}\Pi) = n(\mathcal{L}) - n(\mathcal{L}^{-1}|_{\mathcal{J}^{-1}\ker(\mathcal{L})}). \tag{3.10}$$

It is clearly the case that

$$n(\mathcal{L}) = n(\mathcal{A}) + n(\mathcal{C}).$$

Furthermore, it is straightforward to verify that

$$\mathcal{L}^{-1}|_{\mathcal{J}^{-1}\ker(\mathcal{L})} = (\mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B})|_{\ker(\mathcal{A})}.$$

The instability index (3.9) can then be rewritten as

$$n(\Pi\mathcal{L}\Pi) = n(\mathcal{A}) + n(\mathcal{C}) - n((\mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B})|_{\ker(\mathcal{A})}). \tag{3.11}$$

Combining (3.9) with (3.11) yields the following theorem:

Theorem 3.1. *Suppose that the operators satisfy the assumption of Lemma 2.1. Further suppose that there is a self-adjoint and positive operator \mathcal{S} such that*

- (a) $X_{\mathcal{S}} := \{u \in X : \langle \mathcal{S}u, Su \rangle < \infty\} \subset X$ is dense
- (b) the operators \mathcal{B}, \mathcal{C} are \mathcal{S} -compact, and the operator \mathcal{A} has an \mathcal{S} -compact resolvent.

Finally, assume that the operator \mathcal{C} is invertible. If

- (i) $\mathcal{B}|_{\ker(\mathcal{A})} = \mathbf{0}_{z(\mathcal{A})}$, and
- (ii) $(\mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B})|_{\ker(\mathcal{A})}$ is invertible,

then the total number of eigenvalues in the closed right-half of the complex plane satisfies the instability index

$$k_r + k_c + k_i^- = n(\mathcal{A}) + n(\mathcal{C}) - n((\mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B})|_{\ker(\mathcal{A})}).$$

Remark 3.2. If \mathcal{A} is nonsingular, then Theorem 3.1 is precisely the result of [34, Corollary 3.9]. On the other hand, in the event that \mathcal{A} has a nontrivial kernel then it is an improvement over [34, Theorem 4.2], where it was shown that

$$k_r + k_c + k_i^- \leq n(\mathcal{A}) + n(\mathcal{C}).$$

The proof presented here is quite different than that given in [34]; in particular, in that paper the analysis takes place on Pontryagin spaces, and the linearization that is studied there is not the Hamiltonian linearization of (3.2).

Remark 3.3. If the imaginary part of all of the operators is zero, then due to the Hamiltonian eigenvalue symmetry $\{\pm\lambda, \pm\bar{\lambda}\} \subset \sigma(\mathcal{P}_2)$ it is necessarily the case that k_c and k_i^- are even-value. Thus, under this assumption $n(\Pi\mathcal{L}\Pi)$ being odd automatically implies that $k_r \geq 1$. On the other hand, if one or more of the operators has a nontrivial imaginary part, then the Hamiltonian eigenvalue symmetry reduces to $\{\lambda, -\bar{\lambda}\} \subset \sigma(\mathcal{P}_2)$, and no such conclusion can be drawn.

Remark 3.4. One consequence of the index is that all but a finite number of eigenvalues are purely imaginary; furthermore, the purely imaginary eigenvalues have positive Krein signature if the modulus is sufficiently large.

Remark 3.5. The proof of the index formula (3.9) in [18] first required the SCS Basis Lemma, in which it was shown that the generalized eigenvectors associated with the linearization (3.1) formed a basis. Technical assumptions on the operators were needed in order for the SCS Basis Lemma to hold true. Unfortunately, at least in the application discussed later in this paper these technical assumptions do not hold; hence, the alternate proof via the limiting argument.

4. Applications to Second-Order in Time Hamiltonian Systems

As discussed in the introduction, quadratic operator pencils arise naturally when one studies the stability of solutions to second-order in time Hamiltonian systems. In this section, we present two applications of the general theory developed in the previous sections in precisely this context. We begin by considering the stability of periodic waves in an (abstract) nonlinear wave equation posed in a Hilbert space, and conclude with a stability analysis for periodic waves in the so-called “good” Boussinesq equation.

4.1. *Example: stability in (abstract) nonlinear wave equations.* One important example of quadratic pencils arises in the study of second-order (in time) Hamiltonian systems (for a specific case of the following discussion, see, e.g., [14, Section 7]). Consider a wave equation of the form

$$\partial_t^2 u + \mathcal{H}'(u) = 0, \quad \mathcal{H}^{(k)}(u) := \frac{\delta^k \mathcal{H}}{\delta u^k}(u), \tag{4.1}$$

where $u \in X$, which is a Hilbert space with inner-product $\langle \cdot, \cdot \rangle$. The Hamiltonian $\mathcal{H} : X \mapsto \mathbb{R}$ is assumed to be smooth.

It will be assumed that the Hamiltonian system has symmetries. Let G be a finite-dimensional abelian Lie group with Lie algebra \mathfrak{g} . Denote by $\exp(\omega) = e^\omega$ for $\omega \in \mathfrak{g}$ the exponential map from \mathfrak{g} into G , and assume that $\mathcal{T} : G \mapsto L(V)$, where $X \subset V \subset X^*$ (the dual space of X), is a unitary representation of G on V . It is then the case that $\mathcal{T}'(e)$ maps \mathfrak{g} into the space of closed skew-symmetric operators on V with domain X . The notation $\mathcal{T}_\omega := \mathcal{T}'(e)\omega$ for $\omega \in \mathfrak{g}$ will be used to denote the linear skew-symmetric operator which is the generator of the semigroup $\mathcal{T}(e^{\omega t})$. Using this notation the symmetry assumption becomes that the Hamiltonian satisfies $\mathcal{T}(\omega)\mathcal{H}(u) = \mathcal{H}(\mathcal{T}(\omega)u)$ for all $\omega \in \mathfrak{g}$.

Writing $\mathbf{u} = (u, v)^T$, where $v = \partial_t u \in X_1$ (in applications, it is often the case that $X \subset X_1$ is dense), the system (4.1) can be written on $X \times X_1$ as the first-order Hamiltonian system

$$\partial_t \mathbf{u} = \mathcal{J} \widehat{\mathcal{H}}'(\mathbf{u}), \tag{4.2}$$

where

$$\mathcal{J} = \begin{pmatrix} 0 & \mathcal{I} \\ -\mathcal{I} & 0 \end{pmatrix}, \quad \widehat{\mathcal{H}}(\mathbf{u}) = \mathcal{H}(u) + \frac{1}{2} \langle v, v \rangle.$$

The system (4.2) is invariant under the action $\widehat{\mathcal{T}}(\omega)$, where

$$\widehat{\mathcal{T}}(\omega)\mathbf{u} = \begin{pmatrix} \mathcal{T}(\omega) & 0 \\ 0 & \mathcal{T}(\omega) \end{pmatrix} \mathbf{u}.$$

An n -parameter family of conserved quantities for the Hamiltonian system (4.2) is induced from the self-adjoint operator $\mathcal{J}^{-1}\widehat{\mathcal{T}}_\omega$, and is given by

$$\mathcal{Q}(\mathbf{u}) := \frac{1}{2} \langle \mathcal{J}^{-1}\widehat{\mathcal{T}}_\omega \mathbf{u}, \mathbf{u} \rangle = -\operatorname{Re} \langle \mathcal{T}_\omega u, v \rangle.$$

Upon defining the Lagrangian

$$\Lambda(\mathbf{u}) := \widehat{\mathcal{H}}(\mathbf{u}) + \mathcal{Q}(\mathbf{u}),$$

waves to (4.2) will be realized as steady-state solutions for the system

$$\partial_t \mathbf{u} = \mathcal{J} \Lambda'(\mathbf{u}), \tag{4.3}$$

i.e., they are critical points for the Lagrangian. Since

$$\Lambda'(\mathbf{u}) = \widehat{\mathcal{H}}'(\mathbf{u}) + \mathcal{J}^{-1}\widehat{\mathcal{T}}_\omega(\mathbf{u}) = \begin{pmatrix} \mathcal{H}'(u) + \mathcal{T}_\omega v \\ v - \mathcal{T}_\omega u \end{pmatrix},$$

critical points are solutions to

$$\mathcal{H}'(u) + \mathcal{T}_\omega^2 u = 0, \quad \mathcal{T}_\omega^2 u := \mathcal{T}_\omega(\mathcal{T}_\omega u). \quad (4.4)$$

It should be noted here that (4.3) is equivalent to the second-order problem

$$\partial_t^2 u + 2\mathcal{T}_\omega \partial_t u + \mathcal{H}'(u) + \mathcal{T}_\omega^2 u = 0. \quad (4.5)$$

Suppose that $u = U$ is a solution to (4.4) (the ω -dependence of the solution is being suppressed here), so that $\mathbf{U} = (U, \mathcal{T}_\omega U)^\top$ is a critical point of the Lagrangian. Indeed, further suppose that there is a nonempty open set $\Omega \subset \mathfrak{g}$ such that the solution is smooth in ω for all $\omega \in \Omega$, and further assume that the isotropy subgroups $\{g \in G : \mathcal{T}(g)U = U\}$ are discrete for all ω . Now consider the spectral and orbital stability of the wave. The linearized problem associated with (4.3) is given by

$$\partial_t \mathbf{u} = \mathcal{J} \mathcal{L} \mathbf{u}, \quad (4.6)$$

where the self-adjoint operator \mathcal{L} is

$$\mathcal{L} := \Lambda''(\mathbf{U}) = \begin{pmatrix} \mathcal{H}''(U) & \mathcal{T}_\omega \\ -\mathcal{T}_\omega & \mathcal{I} \end{pmatrix}.$$

The eigenvalue problem for (4.6) is given by

$$\mathcal{J} \mathcal{L} \mathbf{u} = \lambda \mathbf{u}.$$

This eigenvalue problem is the system

$$-\mathcal{H}''(U)u - \mathcal{T}_\omega v = \lambda v, \quad -\mathcal{T}_\omega u + v = \lambda u,$$

which after substitution is equivalent to the quadratic pencil

$$(\mathcal{H}''(U) + \mathcal{T}_\omega^2 + 2\lambda \mathcal{T}_\omega + \lambda^2 \mathcal{I})u = 0. \quad (4.7)$$

In the notation of (1.4) one has

$$\mathcal{A} = \mathcal{H}''(\phi) + \mathcal{T}_\omega^2, \quad \mathcal{B} = 2\mathcal{T}_\omega, \quad \mathcal{C} = \mathcal{I}.$$

Note that the operators \mathcal{A} , \mathcal{C} are self-adjoint, while the operator \mathcal{B} is skew-symmetric. It is interesting to note that the negative index of \mathcal{L} is discussed in [28, Lemma 1], where it is stated that

$$n(\mathcal{L}) = n(\mathcal{H}''(\phi) + \mathcal{T}_\omega^2).$$

The number of negative directions of \mathcal{L} is precisely the number of negative directions associated with the linearization of (4.4) about $u = U$.

With respect to the spectrum of the pencil (4.7) the result of Theorem 3.1 says the following. The assumptions associated with the symmetries present in the problem imply that

$$\ker(\mathcal{H}''(U) + \mathcal{T}_\omega^2) = \text{span}\{\mathcal{T}_\omega U\},$$

so that $z(\mathcal{H}''(U) + \mathcal{T}_\omega^2) = n$. Furthermore, these assumptions imply that

$$\mathcal{T}_\omega : \ker(\mathcal{H}''(U) + \mathcal{T}_\omega^2) \mapsto \ker(\mathcal{H}''(U) + \mathcal{T}_\omega^2)^\perp,$$

so that the generalized kernel for the pencil has (at least) dimension $2n$. Since $\mathcal{C} = \mathcal{I}$, under the assumption that the matrix

$$(\mathcal{I} - 4\mathcal{T}_\omega(\mathcal{H}''(U) + \mathcal{T}_\omega^2)^{-1}\mathcal{T}_\omega)|_{\text{span}\{\mathcal{T}_\omega U\}}$$

is invertible, it will then be the case that the instability index count satisfies

$$k_r + k_c + k_1^- = n(\mathcal{H}''(\phi) + \mathcal{T}_\omega^2) - n\left((\mathcal{I} - 4\mathcal{T}_\omega(\mathcal{H}''(U) + \mathcal{T}_\omega^2)^{-1}\mathcal{T}_\omega)|_{\text{span}\{\mathcal{T}_\omega U\}}\right) \quad (4.8)$$

Now that the spectral problem is understood, consider the orbital stability of the wave. This result follows almost immediately for [14, Theorem 4.1]. An alternate interpretation of that result is as follows. In the language of that paper the wave is said to be orbitally stable if the reduced Hamiltonian, which is the Hamiltonian restricted to the closed subspace orthogonal to the generalized kernel of $\mathcal{J}\mathcal{L}$, is positive definite. As was discussed in, e.g., [8,9,21], this condition is equivalent to saying that for the linearized problem (4.6) the spectrum is purely imaginary and satisfies $k_1^- = 0$. Under this spectral assumption, and the compactness assumptions associated with the operators, the wave is then a local minimizer for the Lagrangian, and hence is orbitally stable.

Theorem 4.1. *Suppose that for the quadratic pencil*

$$\mathcal{P}_2(\lambda) := (\mathcal{H}''(U) + \mathcal{T}_\omega^2) + \lambda(2\mathcal{T}_\omega) + \lambda^2\mathcal{I},$$

which is the spectral problem for the linearization of the second-order Hamiltonian system (4.5) about the steady-state $u = U$, the operators satisfy the assumptions associated with Theorem 3.1. Assume that solutions to (4.5) exist globally in time. If the eigenvalues satisfy the instability index count $k_r = k_1^- = k_c = 0$ (see (4.8)), then the wave is orbitally stable. In other words, for each $\epsilon > 0$ there is a $\delta > 0$ such that if

$$\|u(0) - U\|_X + \|\partial_t u(0) - \mathcal{T}_\omega U\|_{X_1} < \delta,$$

then

$$\sup_{t>0} \inf_{g \in G} (\|u(t) - \mathcal{T}(g)U\|_X + \|\partial_t u(t) - \mathcal{T}(g)\mathcal{T}_\omega U\|_{X_1}) < \epsilon.$$

4.2. *Example: periodic waves to the “good” Boussinesq equation.* The generalized “good” Boussinesq equation (gB) is of the form

$$\partial_t^2 u + \partial_x^2(\partial_x^2 u - u + f(u)) = 0, \quad (4.9)$$

where $f : \mathbb{R} \mapsto \mathbb{R}$ is smooth. In traveling coordinates, i.e., $\xi = x - ct$ with $c \in (-1, 1)$, the gB can be rewritten as

$$\partial_t^2 u - 2c\partial_{t\xi}^2 u + \partial_\xi^2(\partial_\xi^2 u - (1 - c^2)u + f(u)) = 0. \quad (4.10)$$

The interest will be on solutions to (4.10) which are $2L$ -periodic in ξ , i.e., $u(\xi + 2L, t) = u(\xi, t)$.

In order to study the existence, spectral, and orbital stability problems, it is convenient to recast the gB (4.9) in a Hamiltonian formulation similar to that of (4.2). Herein this task will be accomplished via a trick presented in [4]. The evolution is considered to take

place on the space $L^2_{\text{per}}[-L, +L]$, i.e., the space of square-integrable functions which are $2L$ -periodic in ξ . The inner-product is the standard one, i.e.,

$$\langle f, g \rangle = \int_{-L}^{+L} f(x)\overline{g(x)} \, dx.$$

It is straightforward to check that the original gB (4.9) is equivalent to the system

$$\partial_t \mathbf{u} = \mathcal{J} \widehat{\mathcal{H}}'(\mathbf{u}), \quad \mathbf{u} = (u, v)^T, \tag{4.11}$$

where $\partial_x v = \partial_t u$,

$$\mathcal{J} = \begin{pmatrix} 0 & \partial_x \\ \partial_x & 0 \end{pmatrix}, \quad \widehat{\mathcal{H}}(\mathbf{u}) = \int_{-L}^{+L} \left[\frac{1}{2}(\partial_x u)^2 + \frac{1}{2}u^2 - F(u) + \frac{1}{2}v^2 \right] \, dx.$$

Here $F'(u) = f(u)$. Note that the above formulation of $\widehat{\mathcal{H}}$ is consistent with the formulation of the previous section, i.e.,

$$\widehat{\mathcal{H}}(\mathbf{u}) = \mathcal{H}(u) + \frac{1}{2}\langle v, v \rangle, \quad \mathcal{H}(u) = \int_{-L}^{+L} \left[\frac{1}{2}(\partial_x u)^2 + \frac{1}{2}u^2 - F(u) \right] \, dx,$$

while the skew-symmetric operator \mathcal{J} no longer has the property of having a bounded inverse. The system is invariant under spatial translation, i.e., $\widehat{\mathcal{T}}(\omega)\mathbf{u}(x, t) = \mathbf{u}(x + \omega, t)$. Consequently, upon using the fact that on $\ker(\partial_x)^\perp$ it is true that

$$\mathcal{J}^{-1}\widehat{\mathcal{T}}_\omega = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

the conserved quantity associated with the spatial translation is given by

$$\mathcal{Q}(\mathbf{u}) = \langle u, v \rangle \quad \left(= \frac{1}{2}\partial_t \langle u, u \rangle \right),$$

and the Lagrangian for the system is

$$\Lambda(\mathbf{u}) = \widehat{\mathcal{H}}(\mathbf{u}) + c\mathcal{Q}(\mathbf{u}) \quad \Rightarrow \quad \Lambda'(\mathbf{u}) = \begin{pmatrix} \mathcal{H}'(u) + cv \\ cu + v \end{pmatrix}.$$

In conclusion, the system to be studied is

$$\partial_t \mathbf{u} = \mathcal{J} \Lambda'(\mathbf{u}), \tag{4.12}$$

which is equivalent to $\partial_t \mathbf{u} = \widehat{\mathcal{H}}'(\mathbf{u})$ in traveling coordinates $\xi = x - ct$.

First consider the existence problem. Since $\ker(\partial_x) = \text{span}\{1\}$, for real-valued parameters a, b the problem is

$$-\partial_x^2 u + u - f(u) + cv = -a, \quad v + cu = b \tag{4.13}$$

which is equivalent to

$$-\partial_x^2 u + (1 - c^2)u - f(u) = -(a + cb). \tag{4.14}$$




Fig. 1. (color online) The criteria that the potential $V(U, a, \hat{c})$ must satisfy relative to the energy E in order for there to exist spatially periodic solutions. Here the energy was chosen so that there exist two distinct periodic solutions. Increasing the energy past a threshold E^* , for which there are two homoclinic orbits, yields that the two solutions merge into one periodic solution

This is a well-studied problem. In order to use the desired geometric formulation of Bronski et al. [6], it will first be necessary to rescale the wave-speed via

$$\hat{c} := 1 - c^2 \quad \Rightarrow \quad c = c_{\pm} := \pm\sqrt{1 - \hat{c}}, \tag{4.15}$$

so that (4.14) can be rewritten as

$$-\partial_x^2 u + \hat{c}u - f(u) = -(a + cb). \tag{4.16}$$

Note that in (4.16) the wave-speed c is either of c_{\pm} . Without loss of generality assume that $b = 0$. A periodic steady-state, say U , will be a solution to the ODE

$$\partial_{\xi}^2 U - \hat{c}U + f(U) = a; \tag{4.17}$$

hence, the solution will naturally depend on the parameters a and \hat{c} . If one sets

$$E = \frac{1}{2}(\partial_{\xi} U)^2 + V(U, a, \hat{c}), \quad V(U, a, \hat{c}) := -aU - \frac{1}{2}\hat{c}U^2 + F(U),$$

then under the assumption that E , a , and \hat{c} are chosen so that

- (a) $E = V(U, a, \hat{c})$ has (at least) two real roots U_{\pm} with $U_- < U_+$
- (b) $V(U, a, \hat{c}) < E$ for $U_- < U < U_+$

(see Fig. 1) there will be a periodic solution with period $2L$, where

$$L = \frac{1}{\sqrt{2}} \int_{U_-}^{U_+} \frac{dU}{\sqrt{E - V(U, a, \hat{c})}}.$$

As it will be seen, in particular examples the spectral stability of the $2L$ -periodic solution will naturally depend upon the parameters a, \hat{c}, E . The dependence of the solution on these parameters will be implicit in all that follows.

Now consider the stability problem. Let the $2L$ -periodic wave found in (4.16) be denoted by $u = U$. Recalling that we set $b = 0$, from (4.13) the v -component of the wave is given by $v = -cU$, where from (4.15) one has $c = c_{\pm}$ can be of either sign for a fixed value of \hat{c} . The steady-state solution for (4.12) is then given by

$$\mathbf{u} = U = \begin{pmatrix} U \\ -cU \end{pmatrix}.$$

Under the mapping

$$u = U + v \tag{4.18}$$

the system (4.12) becomes

$$\partial_t v = \mathcal{J} \Lambda'(U + v).$$

The evolution problem must now be considered on a space for which \mathcal{J} has bounded inverse, i.e., on the space of mean-zero functions. Let $\Pi_0 : L^2_{\text{per}}[-L, +L] \mapsto H_0$ be the self-adjoint projection operator

$$\Pi_0 u = u - \frac{1}{2L} \langle u, 1 \rangle;$$

in other words, Π_0 is the orthogonal projection onto $\ker(\partial_\xi)^\perp$. Here

$$H_0 := \{u \in L^2_{\text{per}}[-L, +L] : \langle u, 1 \rangle = 0\} = \ker(\partial_\xi)^\perp.$$

When writing $\Pi_0 v$ it will be implicitly assumed that Π_0 is being applied to each component of v . In [8, Section 2] it is shown that the proper evolution equation to consider is

$$\partial_t v = \mathcal{J} \Pi_0 \Lambda'(U + v), \quad v(0) = v_0, \tag{4.19}$$

where $\Pi_0 v_0 = v_0$ implies that $\Pi_0 v(t) = v(t)$ for all $t > 0$. In other words, (4.19) describes the evolution of mean-zero perturbations of the underlying wave. Since the evolution occurs on $H_0 \times H_0$, and $\partial_x : H_0 \mapsto H_0$ has bounded inverse, in this formulation the operator \mathcal{J} now has bounded inverse.

First consider the spectral stability problem. The linearized eigenvalue problem

$$\lambda v = \mathcal{J} \Pi_0 \Lambda''(U) v, \quad \Pi_0 v = v$$

can be rewritten as

$$\partial_x [\Pi_0 \mathcal{H}''(U) \Pi_0 u + c v] = \lambda v, \quad \partial_x [c u + v] = \lambda u,$$

where $u, v \in H_0$. Differentiating the first equation yields

$$\partial_x^2 \Pi_0 \mathcal{H}''(U) \Pi_0 u + c \partial_x (\partial_x v) = \lambda \partial_x v,$$

and substituting the second equation into the first and simplifying gives the quadratic pencil problem

$$\left[\lambda^2 - 2c\lambda \partial_x + \partial_x^2 (\Pi_0 (-\mathcal{H}''(U) + c^2) \Pi_0) \right] u = 0.$$

Since $u \in H_0$, one can write $v = \partial_x^{-1} u \in H_0$, so that the pencil becomes

$$\partial_x \left[\lambda^2 - 2c\lambda \partial_x - \partial_x (\Pi_0 (\mathcal{H}''(U) - c^2) \Pi_0) \partial_x \right] v = 0.$$

Since ∂_x has bounded inverse, upon setting \mathcal{L}_2 to be the well-understood self-adjoint Hill operator

$$\mathcal{L}_2 = -\partial_x^2 + \hat{c} - f'(U(x)),$$

the pencil problem to be studied is

$$\left[\lambda^2 - 2c\lambda\partial_x - \partial_x(\Pi_0\mathcal{L}_2\Pi_0)\partial_x \right] v = 0, \quad v \in H_0.$$

Note that in the notation of the previous section,

$$\mathcal{C} = \mathcal{I}, \quad \mathcal{B} = -2c\partial_x, \quad \mathcal{A} = -\partial_x(\Pi_0\mathcal{L}_2\Pi_0)\partial_x. \tag{4.20}$$

Before proceeding with the spectral analysis, the assumptions on the operators given in Theorem 3.1 must be verified. First consider $\ker(\mathcal{A})$. Since $\mathcal{L}_2(\partial_\xi U) = 0$, it is true that

$$\mathcal{L}_2\Pi_0 \cdot \partial_\xi(U - \bar{U}) = 0, \quad \bar{U} = \frac{1}{2L} \int_{-L}^{+L} U(x) dx.$$

In other words, $U - \bar{U} \in \ker(\mathcal{A})$. In order for $\ker(\mathcal{A})$ to have another linearly independent element, it must be the case that $\mathcal{L}_2^{-1}(1) \in H_0$. It will be henceforth assumed that no other element in the kernel exists, i.e.,

$$\langle \mathcal{L}_2^{-1}(1), 1 \rangle \neq 0, \tag{4.21}$$

so that

$$\ker(\mathcal{A}) = \text{span}\{U - \bar{U}\}.$$

Letting

$$P_{\mathcal{A}} : H_0 \mapsto \text{span}\{U - \bar{U}\}, \quad P_{\mathcal{A}}^\perp : H_0 \mapsto \text{span}\{U - \bar{U}\}^\perp \subset H_0$$

be orthogonal projections, it must be checked that

$$(P_{\mathcal{A}}^\perp \mathcal{A} P_{\mathcal{A}}^\perp)^{-1} P_{\mathcal{A}}^\perp \mathcal{B} P_{\mathcal{A}}^\perp, \quad (P_{\mathcal{A}}^\perp \mathcal{A} P_{\mathcal{A}}^\perp)^{-1} P_{\mathcal{A}}^\perp \mathcal{C} P_{\mathcal{A}}^\perp$$

are compact operators. This immediately follows from the fact that $(P_{\mathcal{A}}^\perp \mathcal{A} P_{\mathcal{A}}^\perp)^{-1}$ is compact, and both \mathcal{B} and \mathcal{C} are differentiable operators of lesser order than \mathcal{A} . Together, the above considerations verify the hypothesis of Lemma 2.1.

Next, we must verify that the operators \mathcal{A} , \mathcal{B} , and \mathcal{C} are \mathcal{S} -compact for some compact operator \mathcal{S} . For each $\alpha > 0$, define the operator

$$\mathcal{S}_\alpha := \Pi_0(\partial_x^2 + 1)^\alpha \Pi_0$$

acting on $L^2_{\text{per}}([-L, +L])$. It is clear that \mathcal{S}_α^{-1} is a compact self-adjoint operator on H_0 for each $\alpha > 0$; furthermore, it is true that the space

$$X_{\mathcal{S}_\alpha} = \{u \in H_0 : \langle \Pi_0(\partial_x^2 + 1)^{2\alpha} \Pi_0 u, u \rangle < \infty\}$$

is dense for any $\alpha > 0$. Now, clearly the operator $\mathcal{S}_\alpha^{-1} \mathcal{C} \mathcal{S}_\alpha^{-1} = \mathcal{S}_\alpha^{-2}$ is compact for any $\alpha > 0$. Regarding the operator \mathcal{B} , it is easy to see that $\mathcal{S}_\alpha^{-1} \partial_x \mathcal{S}_\alpha^{-1}$ will be compact as long as $1/4 < \alpha$. Finally, the operator $\mathcal{S}_\alpha^{-1} \mathcal{A} \mathcal{S}_\alpha^{-1}$ will have a compact resolvent as long as $0 < \alpha < 1$. In conclusion, as long as $1/4 < \alpha < 1$, the operators will be \mathcal{S}_α -compact, thus verifying hypothesis (a) and (b) of Theorem 3.1.

From the skew-symmetry of the operator and the fact that U is $2L$ -periodic it is clear that

$$\mathcal{B}|_{\ker(\mathcal{A})} = -2c\langle U - \bar{U}, \partial_x(U - \bar{U}) \rangle = 0.$$

Provided that $(\mathcal{I} - 4c^2\partial_x\mathcal{A}^{-1}\partial_x)|_{\ker(\mathcal{A})}$ is invertible then, a direct application of Theorem 3.1, using the explicit form of the operators given in (4.20) and noting that $\mathcal{C} = \mathcal{I}$ is clearly a positive definite operator, implies that the index count satisfies

$$k_r + k_c + k_i^- = n(\mathcal{A}) - n((\mathcal{I} - 4c^2\partial_x\mathcal{A}^{-1}\partial_x)|_{\ker(\mathcal{A})}).$$

In order to compute $n(\mathcal{A})$, first note that

$$\langle u, \mathcal{A}u \rangle = \langle u, -\partial_x(\Pi_0\mathcal{L}_2\Pi_0)\partial_x u \rangle = \langle \partial_x u, \Pi_0\mathcal{L}_2\Pi_0(\partial_x u) \rangle.$$

Thus, upon using the fact that $\partial_x : H_0 \mapsto H_0$ has a bounded inverse it is clear that

$$n(\mathcal{A}) = n(\Pi_0\mathcal{L}_2\Pi_0).$$

Regarding the quantity on the right, it was shown in [8, equation (2.25)] that if the inequality of (4.21) holds, then

$$n(\Pi_0\mathcal{L}_2\Pi_0) = n(\mathcal{L}_2) - n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle).$$

Consequently, it can now be said that

$$n(\mathcal{A}) = n(\mathcal{L}_2) - n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle), \tag{4.22}$$

so that the index count satisfies

$$k_r + k_c + k_i^- = n(\mathcal{L}_2) - n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle) - n((\mathcal{I} - 4c^2\partial_x\mathcal{A}^{-1}\partial_x)|_{\ker(\mathcal{A})}).$$

Recalling that $c^2 = 1 - \hat{c}$, the index count is complete once the scalar

$$(\mathcal{I} - 4(1 - \hat{c})\partial_x\mathcal{A}^{-1}\partial_x)|_{\ker(\mathcal{A})}$$

is computed. From (4.20) one has that

$$\begin{aligned} \mathcal{I} - 4(1 - \hat{c})\partial_x\mathcal{A}^{-1}\partial_x &= \mathcal{I} + 4(1 - \hat{c})\partial_x \cdot \partial_x^{-1}(\Pi_0\mathcal{L}_2\Pi_0)^{-1}\partial_x^{-1} \cdot \partial_x \\ &= \mathcal{I} + 4(1 - \hat{c})(\Pi_0\mathcal{L}_2\Pi_0)^{-1}; \end{aligned}$$

the second line follows from the fact that ∂_x is invertible on H_0 . Using the characterization of $\ker(\mathcal{A})$, it is then seen that

$$\begin{aligned} &(\mathcal{I} - 4(1 - \hat{c})\partial_x\mathcal{A}^{-1}\partial_x)|_{\ker(\mathcal{A})} \\ &= \langle U - \bar{U}, U - \bar{U} \rangle + 4(1 - \hat{c})\langle (\Pi_0\mathcal{L}_2\Pi_0)^{-1}(U - \bar{U}), U - \bar{U} \rangle. \end{aligned}$$

Finally, in the study of the orbital stability of periodic waves for the generalized Korteweg–de Vries equation (gKdV) it was shown in [8, Section 3] that

$$\langle (\Pi_0\mathcal{L}_2\Pi_0)^{-1}(U - \bar{U}), U - \bar{U} \rangle = D_{\text{gKdV}}, \quad D_{\text{gKdV}} := \frac{\begin{vmatrix} \langle \mathcal{L}_2^{-1}(U), U \rangle & \langle \mathcal{L}_2^{-1}(U), 1 \rangle \\ \langle \mathcal{L}_2^{-1}(U), 1 \rangle & \langle \mathcal{L}_2^{-1}(1), 1 \rangle \end{vmatrix}}{\langle \mathcal{L}_2^{-1}(1), 1 \rangle}.$$

Lemma 4.2. *Consider the quadratic pencil (4.19). If $\langle \mathcal{L}_2^{-1}(1), 1 \rangle \neq 0$, then the stability index is given by*

$$k_r + k_i^- + k_c = n(\mathcal{L}_2) - n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle) - n(\langle U - \bar{U}, U - \bar{U} \rangle) + 4(1 - \hat{c})D_{\text{gKdV}},$$

where

$$D_{\text{gKdV}} := \frac{\begin{vmatrix} \langle \mathcal{L}_2^{-1}(U), U \rangle & \langle \mathcal{L}_2^{-1}(U), 1 \rangle \\ \langle \mathcal{L}_2^{-1}(U), 1 \rangle & \langle \mathcal{L}_2^{-1}(1), 1 \rangle \end{vmatrix}}{\langle \mathcal{L}_2^{-1}(1), 1 \rangle},$$

and the parameter \hat{c} is related to the original wave-speed c via $c^2 = 1 - \hat{c}$.

Remark 4.3. It was shown in [8, Theorem 2.6] that the stability index for the gKdV is given by

$$k_r + k_i^- + k_c = n(\mathcal{L}_2) - n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle) - n(D_{\text{gKdV}});$$

hence, when studying the spectrum for periodic waves to gB and gKdV there is an intimate connection in the indices for the two problems. If $D_{\text{gKdV}} < 0$, then for

$$1 > \hat{c} > 1 + \frac{\langle U - \bar{U}, U - \bar{U} \rangle}{4D_{\text{gKdV}}}$$

the stability index for the gB is exactly the same as for the gKdV. Otherwise, there is precisely one more eigenvalue which is counted by the index. On the other hand, if $D_{\text{gKdV}} > 0$, then the index for the quadratic pencil is exactly that for the gKdV equation.

Remark 4.4. There is a geometric interpretation associated with the quantity D_{gKdV} . The interested reader should consult [6] for more details.

Remark 4.5. In Hakkaev et al. [16] the spectral stability problem was considered under the additional assumption that $n(\mathcal{L}_2) = 1$. Furthermore, while it is not explicitly stated, they further assume that $\langle \mathcal{L}_2^{-1}(1), 1 \rangle > 0$, so that (in this paper’s notation) the index becomes

$$k_r + k_i^- + k_c = 1 - n(\langle U - \bar{U}, U - \bar{U} \rangle) + 4(1 - \hat{c})D_{\text{gKdV}}.$$

The instability criterion in that paper follows from the fact that k_i^-, k_c must be even integers, so that $k_i^- = k_c = 0$, with

$$k_r = \begin{cases} 1, & \hat{c} > \hat{c}^* \\ 0, & \hat{c} < \hat{c}^*, \end{cases}$$

where

$$\hat{c}^* = 1 + \frac{\langle U - \bar{U}, U - \bar{U} \rangle}{4D_{\text{gKdV}}}.$$

It is not clear if in that paper an explicit connection is shown between the gB and the gKdV.

Now consider the orbital stability problem. The local global well-posedness problem has been studied in, e.g., [3, 10, 11], and it will henceforth be assumed that the problem can be solved (at least) locally. Depending on the growth rate of the nonlinearity, this implies that the initial data for (4.19) satisfies, e.g., $v_1(0) \in H_{\text{per}}^1[-L, +L]$, and $v_2(0) \in H_{\text{per}}^{-1}[-L, +L]$, where the norm for the latter space is given by

$$\|u\|_{H^{-1}}^2 = \sum_{z \in \mathbb{Z}} \frac{|\hat{u}(z)|^2}{1 + |z|^2}.$$

The form of the Lagrangian in (4.12) makes clear that in order to control the nonlinear terms the proper space in which to work is $H_{\text{per}}^1[-L, +L] \times L_{\text{per}}^2[-L, +L]$. It will be further assumed that the hypothesis leading to Lemma 4.2 hold, and that the spectral problem has zero instability index, i.e., $k_r = k_c = k_i^- = 0$. As was seen in [8, Section 2.4], this is sufficient in order to conclude that the wave is orbitally stable with respect to the evolution defined by (4.19).

Proposition 4.6. *Suppose that the IVP for (4.19) is locally well-posed. Further suppose that in addition to what is required for a unique local solution to exist, the mean-free perturbative initial data for the system (4.19) satisfies $\mathbf{v}(0) \in H_{\text{per}}^1[-L, +L] \times L_{\text{per}}^2[-L, +L]$. If the spectral problem satisfies $k_r = k_c = k_i^- = 0$, then the underlying wave is orbitally stable. In other words, for each $\epsilon > 0$ there is a $\delta > 0$ such that for (4.19),*

$$\|\mathbf{u}(0) - \mathbf{U}\|_{H_{\text{per}}^1 \times L_{\text{per}}^2} < \delta \implies \sup_{t > 0} \inf_{\omega \in \mathbb{R}} \|\mathbf{u}(t) - \widehat{\mathcal{T}}(\omega)\mathbf{U}\|_{H_{\text{per}}^1 \times L_{\text{per}}^2} < \epsilon.$$

Note that for the original system (4.9) the requirement on the initial data is

$$u(0) = U + v_1(0), \quad \partial_t u(0) = -c\partial_x U + \partial_x v_2(0),$$

where each component $v_j(0)$ has zero mean. A natural question is then: what happens if the initial perturbation is not mean-free? In this case, we now argue for orbital stability with respect to a nearby periodic traveling wave of (4.19). For related arguments in the contexts of other nonlinear dispersive equations, see the work of Hărăgus and Gallay [17] on the nonlinear Schrodinger equation as well as the works of Johnson [19, 20] on generalized KdV and BBM models, respectively.

To begin, we make a few comments regarding the conserved quantities of (4.11). As discussed above, even though we set $b = 0$ in the analysis, the periodic traveling wave solutions of (4.19), which are solutions to the ODE (4.14), form a five-parameter family of solutions of the form

$$u_{\xi}(x, t) = u(x - ct + \xi; a, E, c, b)$$

where $\xi \in \mathbb{R}$ and (a, E, c, b) belong to some open set $\Omega \in \mathbb{R}^4$. Furthermore, the evolution equation (4.11) admits the following two conserved quantities: the momentum (charge)

$$\mathcal{P}(\mathbf{w}) := \int_{-L}^L w_1 w_2 \, dx, \quad \mathbf{w} := (w_1, w_2) \in H_{\text{per}}^1[-L, +L] \times L_{\text{per}}^2[-L, +L],$$

arising from the translation invariance of (4.19), and the casimirs

$$\begin{aligned} \mathcal{M}_1(\mathbf{w}) &:= \int_{-L}^L w_1 \, dx, \quad \mathcal{M}_2(\mathbf{w}) := \int_{-L}^L w_2 \, dx, \\ \mathbf{w} &:= (w_1, w_2) \in H_{\text{per}}^1[-L, +L] \times L_{\text{per}}^2[-L, +L], \end{aligned}$$

arising from the fact that $\ker(\mathcal{J})$ is non-trivial. Notice that \mathcal{P} and \mathcal{M} are smooth functionals on $H_{\text{per}}^1[-L, +L] \times L_{\text{per}}^2[-L, +L]$ and that, when restricted to the manifold of traveling wave solutions of (4.19) the functionals \mathcal{M}_1 , \mathcal{M}_2 , and \mathcal{P} reduce to

$$M_1(a, E, c, b) := \int_0^T u(x; a, E, c, b) \, dx, \quad M_2(a, E, c, b) := cM_1(a, E, c, b) - bT,$$

and

$$\tilde{P}(a, E, c, b) := -P(a, E, c, b) + bM_1(a, E, c, b),$$

where here $T = T(a, E, c, b) (= 2L)$ denotes the period of the wave and $P(a, E, c, b) := c \int_0^T u(x; a, E, c, b)^2 \, dx$.

Now, consider the case where the means of $v_j(0)$ are small, but non-zero. Using the geometric formalism of Bronski et al. (see [5,6]) we have the following key lemma.

Lemma 4.7. *With the notation as above, we the equality*

$$\left(\mathcal{I} - 4(1 - \hat{c})\partial_x \mathcal{A}^{-1} \partial_x \right) |_{\ker(\mathcal{A})} = T \det \begin{pmatrix} T_a & M_{1,a} \\ T_E & M_{1,E} \end{pmatrix} \det \begin{pmatrix} T_E & M_{1,E} & \tilde{P}_E & M_{2,E} \\ T_a & M_{1,a} & \tilde{P}_a & M_{2,a} \\ T_c & M_{1,c} & \tilde{P}_c & M_{2,c} \\ T_b & M_{1,b} & \tilde{P}_b & M_{2,b} \end{pmatrix}.$$

Proof. To compute the left hand side, define the function

$$\phi_2(x) := \det \begin{pmatrix} u_a & T_a & M_{1,a} \\ u_E & T_E & M_{1,E} \\ u_c & T_c & M_{1,c} \end{pmatrix}$$

and satisfies

$$\Pi_0 \mathcal{L}_2 \Pi_0 \phi_2 = \Pi_0 \mathcal{L}_2 \phi_2 = 2c \det \begin{pmatrix} T_a & M_{1,a} \\ T_E & M_{1,E} \end{pmatrix} (U - \bar{U}).$$

The stated equality now follows by directly calculating

$$\left(\mathcal{I} - 4(1 - \hat{c})\partial_x \mathcal{A}^{-1} \partial_x \right) |_{\ker(\mathcal{A})} = \left\langle \Pi_0 U, \left(1 + 4c^2 (\Pi_0 \mathcal{L}_2 \Pi_0)^{-1} \right) \Pi_0 U \right\rangle$$

and comparing to the right hand side of the above equality. \square

From Lemma 4.7 and the assumption that

$$\left(\mathcal{I} - 4(1 - \hat{c})\partial_x \mathcal{A}^{-1} \partial_x \right) |_{\ker(\mathcal{A})}$$

is nonsingular at the underlying wave U , corresponding say to $(a, E, c, b) = (a_0, E_0, c_0, b_0)$, implies that the map

$$\mathbb{R}^4 \ni (a, E, c, b) \mapsto (T(a, E, c, b), M_1(a, E, c, b), \tilde{P}(a, E, c, b), M_2(a, E, c, b)) \in \mathbb{R}^4$$

is a local diffeomorphism from a neighborhood of (a_0, E_0, c_0, b_0) onto a neighborhood of the point

$$(T, M_1, \tilde{P}, M_2)(a_0, E_0, c_0, b_0).$$

It follows that we can find a curve $[0, 1] \ni s \mapsto (a(s), E(s), c(s), b(s)) \in \mathbb{R}^4$ with $(a(0), E(0), c(0), b(0)) = (0, 0, 0, 0)$ such that for each $s \in [0, 1]$ the function

$$\tilde{u}(x; s) = u(x; a_0 + a(s), E_0 + E(s), c_0 + c(s), b_0 + b(s))$$

is a $T = T(a_0, E_0, c_0, b_0)$ -periodic traveling wave solution of (4.9) and that, moreover the endpoint condition

$$\begin{aligned} M_j(a_0 + a(1), E_0 + E(1), c_0 + c(1), b_0 + b(1)) &= \mathcal{M}_j(\mathbf{u}(0) + \mathbf{v}(0)), \quad j = 1, 2 \\ P(a_0 + a(1), E_0 + E(1), c_0 + c(1), b_0 + b(1)) &= \mathcal{P}(\mathbf{u}(0) + \mathbf{v}(0)) \end{aligned}$$

and growth constraint

$$\sup_{s \in (0, 1)} |(a(s), E(s), c(s), b(s))|_{\mathbb{R}^4} \lesssim \|\mathbf{v}(0)\|_{H^1_{\text{per}}(-L, L) \times L^2_{\text{per}}(-L, L)}$$

are satisfied. Assuming $\|\mathbf{v}(0)\|_{H^1_{\text{per}} \times L^2_{\text{per}}}$ is sufficiently small, it follows that the wave $\tilde{u}(\cdot, 1)$ is nonlinearly orbitally stable in the sense described in Theorem 4.6, which, by the triangle inequality, implies orbital stability of U to initial perturbations $\mathbf{v}(0)$ with nonzero, but sufficiently small, mean. Since \mathcal{M}_j and \mathcal{P} are continuous in the $H^1_{\text{per}} \times L^2_{\text{per}}$ topology, it follows that we have orbital stability in the standard sense without the restriction to mean-free initial data in (4.19). This observation yields the following extension of Theorem 4.6

Theorem 4.8. *Suppose that the IVP for (4.19) is locally well-posed. Further, suppose that in addition to what is required for a unique local solution to exist, the perturbative initial data for the system (4.19) satisfies $\mathbf{v}(0) \in H^1_{\text{per}}[-L, +L] \times L^2_{\text{per}}[-L, +L]$. If the spectral problem satisfies $k_r = k_c = k_i^- = 0$, then the underlying wave is orbitally stable, i.e., for each $\epsilon > 0$ there exists a $\delta > 0$ such that for (4.19) we have*

$$\|\mathbf{u}(0) - U\|_{H^1_{\text{per}} \times L^2_{\text{per}}} < \delta \quad \Rightarrow \quad \sup_{t > 0} \inf_{\omega \in \mathbb{R}} \|\mathbf{u}(t) - \widehat{T}(\omega)U\|_{H^1_{\text{per}} \times L^2_{\text{per}}} < \epsilon.$$

In the next sections, we utilize the above explicit connection of the stability problems for gB and gKdV type equations to make several comments regarding the stability of periodic waves in the gB equation.

4.2.1. *Case study: power law nonlinearity* The stability indices for the gKdV equation were computed in Bronski et al. [6] for the case that $f(u) = u^{p+1}$ for some $p \geq 1$ and $|\hat{c}| < 1$. Note via (4.15) that this implies $|c_{\pm}| < 1$. While we will not do it here, the gB can be rescaled so that the influence of \hat{c} is removed from the steady-state problem. This independence is reflected in the existence diagrams. As in the analysis leading to the statement of Lemma 4.2, it will be assumed here that $b = 0$.

First suppose that $p = 1$, which corresponds to the classical gB equation. In [6, Section 5.1] it is shown that all periodic waves in this model satisfy

$$n(\mathcal{L}_2) = 1, \quad n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle) = 0, \quad n(D_{\text{gKdV}}) = 1.$$

Thus, via Lemma 4.2 it is true that for a given (a, E) there is a critical positive wave-speed $\hat{c}_{a,E}^* < 1$ such that $k_1^- = k_c = 0$ with

$$k_r = \begin{cases} 1, & \hat{c} > \hat{c}_{a,E} \\ 0, & \hat{c} < \hat{c}_{a,E}. \end{cases}$$

Returning to the original wavespeed c , it follows that for any periodic traveling wave solution of the classical gB equation there exists a range of wavespeeds $(1 - \hat{c}_{a,E}^*)^{1/2} < |c| < 1$ for which the wave is nonlinearly (orbitally) stable, while it is unstable spectrally unstable to perturbations with the same period if $|c| < (1 - \hat{c}_{a,E}^*)^{1/2}$. This is consistent with the result of [16, Theorem 2], where $\hat{c}_{a,E}$ is explicitly given when $a = 0$. In that paper the case of nonzero a was not considered and only spectral instability for $|c| \leq (1 - \hat{c}_{a,E}^*)^{1/2}$ was verified. Here, our calculations complemented this result by verifying that waves traveling with speed greater than $(1 - \hat{c}_{a,E}^*)^{1/2}$ are by Theorem 4.6 indeed nonlinearly stable.

For examples which are not covered in Hakkaev et al. [16], e.g., when it is possible for $n(\mathcal{L}_2) \geq 2$, first consider the problem when $p = 2$. The table below, which corresponds to Fig. 2, can be derived from [6, Section 5.2]:

Region	$n(\mathcal{L}_2)$	$n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle)$	$n(D_{\text{gKdV}})$
(a)	1	0	1
(b)	2	0	1
(c)	2	1	0
(d)	2	1	1
(e)	1	0	1

From the theoretical result in Lemma 4.2 it will be the case that in that in regions (a), (d), and (e) there will exist a $0 < \hat{c}_{a,E} < 1$ such that $k_r = 1$ for $\hat{c} > \hat{c}_{a,E}$ and $k_r = 0$ otherwise; furthermore, it is always true that $k_1^- = k_c = 0$ in these regions. In region (c) it will be the case that $k_r = 1$ for all c with the other two indices being zero. Finally, in region (b) there will exist a $0 < \hat{c}_{a,E} < 1$ such that $k_r = 1$ for $-1 < \hat{c} < \hat{c}_{a,E}$ with the other two indices being zero, while for $\hat{c} > \hat{c}_{a,E}$ all that can be said is that $k_r + k_1^- + k_c = 2$. Notice, however, that by parity we see for speeds $\hat{c} > \hat{c}_{a,E}^*$ in region (b) we have $k_r = 0$ and $k_1^- + k_c = 2$, which allows the possibility that some waves may still be spectrally stable in this region with $k_1^- = 2$ or that some waves may be spectrally unstable to perturbations with the same period with $k_c = 2$: such a situation is precluded in the well-studied solitary wave theory.




Fig. 2. (color online) The configuration space in the aE -plane when $p = 2, b = 0$, and $|\hat{c}| < 1$ fixed (see [6, Figure 3]). The swallowtail figure divides the plane into regions containing 0, 1, and 2 periodic solutions. In region (a) there are two solutions, while all of the other marked regions have one solution. In the unmarked region there are no periodic solutions. The quantities used in the stability calculation are given in an accompanying table

Remark 4.9. In [16, Theorem 1] it is shown that for one of the two solutions in region (a) of Fig. 2 with $a = 0$ the index satisfies $k_1^- = k_c = 0$ with $k_r = 1$ for $\hat{c} > \hat{c}_{a,E}$, and $k_r = 0$ for $\hat{c} < \hat{c}_{a,E}$. Furthermore, for this solution the constant $\hat{c}_{a,E}$ is explicitly given when $a = 0$. Although they do not consider the case in their paper, the same result holds in region (e). The parameter region which is outside their theory comprises the union of (b), (c), and (d).

The below table for $p = 4$ which corresponds to Fig. 3 can be derived from [6, Section 5.3]:

Region	$n(\mathcal{L}_2)$	$n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle)$	$n(D_{\text{gKdV}})$
(a)	1	0	1
(a')	1	0	1
(b)	1	0	0
(c)	2	1	0
(d)	2	0	1

From the theoretical result in Lemma 4.2 it will be the case that in that in regions (a) and (a') there will exist a $0 < c_{a,E}^* < 1$ such that $k_r = 1$ for $\hat{c} > \hat{c}_{a,E}$ and $k_r = 0$ otherwise; furthermore, it is always true that $k_1^- = k_c = 0$. In regions (b) and (c) it will be the case that $k_r = 1$ for all \hat{c} with the other two indices being zero. Finally, in region (d) there will exist a $0 < \hat{c}_{a,E} < 1$ such that $k_r = 1$ for $\hat{c} < \hat{c}_{a,E}$ with the other two indices being zero, while for $\hat{c} > \hat{c}_{a,E}$ all that can be said is that $k_r + k_1^- + k_c = 2$.




Fig. 3. (color online) The configuration space in the aE -plane when $p = 4$ and $|\hat{c}| < 1$ (see [6, Figure 4]). The swallowtail figure divides the plane into regions containing 0, 1, and 2 periodic solutions. In region (a) there are two solutions, while all of the other marked regions have one solution. In the unmarked region there are no periodic solutions. The quantities used in the stability calculation are given in an accompanying table

4.2.2. Case study, continued: solitary wave and equilibrium solution limits In this final section, we make some comments regarding the stability of periodic traveling wave solutions of (4.9) which are either near the solitary wave or near an equilibrium (constant) solution. Throughout this section, we continue to consider (4.9) with power law nonlinearity $f(u) = u^{p+1}$ for some $p \geq 1$. In this case, we have from (4.14) that the profile U satisfies the ODE

$$\partial_x^2 u = (1 - c^2)u - u^{p+1},$$

where, for simplicity, we are restricting our discussion to those waves with $a = b = 0$.¹ This equation is clearly Hamiltonian, and has critical points $(u, \partial_x u) = (0, 0)$, corresponding to a saddle point, and $(u, \partial_x u) = ((1 - c^2)^{1/p}, 0)$,² corresponding to a nonlinear center. Further, for a fixed wavespeed $c \in (-1, 1)$, in the two-dimensional $(u, \partial_x u)$ phase plane the nonlinear center $((1 - c^2)^{1/p}, 0)$ is surrounded by a one parameter family of periodic orbits, which are in turn bounded by an orbit which is homoclinic to the saddle point $(0, 0)$. These periodic orbits can be parameterized by the ODE energy E determined from the defining relation

$$\frac{1}{2}(\partial_x u)^2 = E + \frac{1 - c^2}{2}u^2 - \frac{1}{p + 2}u^{p+2}.$$

¹ As we will see below, this is a natural restriction when considering the limiting case to a solitary wave asymptotic to zero as $x \rightarrow \pm\infty$.

² Notice when p is an even integer, the point $(u, \partial_x u) = (-(1 - c^2)^{1/p}, 0)$ is also a critical point. In this discussion, we ignore this additional critical point, noting that any conclusions for periodic waves emerging from the $((1 - c^2)^{1/p}, 0)$ critical point hold also for those emerging from the $(-(1 - c^2)^{1/p}, 0)$ critical point. For general $p \geq 1$, the governing ODE does not admit such negative solutions.

The period $T(E, c)$ of these waves inside the homoclinic orbit satisfies

$$\lim_{E \rightarrow 0^-} T(E, c) = +\infty, \quad \lim_{E \rightarrow E^*(c)^+} T(E, c) = \frac{2\pi}{\sqrt{(1 - c^2)p}}$$

where $E^*(c)$ is the ODE energy level associated with the equilibrium point $((1 - c^2)^{1/p}, 0)$. Below, we consider the stability of the periodic traveling wave solutions of (4.9) in both the distinguished limits $E \rightarrow 0^-$, corresponding to the solitary wave limit, and $E \rightarrow E^*(c)^+$, associated with small amplitude periodic wave trains.

We begin by considering the stability of the periodic traveling waves of (4.9) in the solitary wave limit. When considering solitary wave solutions which decay to zero as $x \rightarrow \pm\infty$ we obtain a two-parameter family of solutions parameterized by wavespeed c and spatial translation, which corresponds to a one-parameter family of homoclinic orbits in the two-dimensional phase space of (4.17) (notice the boundary conditions in this case require $a = E = b = 0$). Keeping throughout $a = b = 0$ and fixing the wavespeed $c \in (-1, 1)$, by the above considerations there exists, up to translations, a one parameter family of “large period” periodic traveling wave solutions of (4.9) parameterized by the ODE energy E . Furthermore, for fixed c these periodic waves approach locally uniformly on \mathbb{R} an appropriate translate of the limiting solitary wave as $E \rightarrow E^-$. The stability of the limiting solitary waves was investigated by Bona and Sachs [4], where they applied the abstract theory of Grillakis et al. [13] to obtain nonlinear stability of the solitary wave when $1 < p < 4$ and $p/4 < c^2 < 1$. This stability theory was later complemented by Liu [29], obtaining nonlinear instability if either $c^2 \leq p/4$ or $p \geq 4$. Next, we show that this phenomenon is observed again for the “nearby” periodic traveling wave solutions of (4.9). For such periodic waves, it is easy to see that

$$n(\mathcal{L}_2) - n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle) = 1, \quad 0 < -E \ll 1,$$

and, furthermore,

$$\lim_{\tilde{E} \rightarrow 0^-} D_{\text{gKdV}} \Big|_{(a,E,c,b)=(0,\tilde{E},c,0)} = -\Lambda(c, p)(4 - p)$$

for some positive constant $\Lambda(c, p) > 0$ (for details, see the asymptotic analysis in [5, Section 3.2]). It follows that for all³ $1 \leq p < 4$ there exists a critical wavespeed $c = c^*(p, E)$ such that the periodic traveling wave $u(\cdot; a = 0, E, c, b = 0)$ with $0 < -E \ll 1$ is orbitally stable for $c^*(p, E) < |c| < 1$ and spectrally unstable with $k_r = 1$ for $|c| < c^*(p, E)$, while for $p > 4$ all such long-period waves are spectrally unstable with $k_r = 1$. This is consistent with the solitary wave orbital stability/instability results of [4] and [29].

Finally, continuing to restrict to $a = b = 0$, we consider the stability of small amplitude periodic traveling waves of (4.9) associated to those periodic orbits of the profile ODE near the nonlinear center $((1 - c^2)^{1/p}, 0)$. Fixing the wavespeed $c \in (-1, 1)$, it follows by basic asymptotic analysis near the equilibrium solution $u = (1 - c^2)^{1/p}$ that

$$n(\mathcal{L}_2) - n(\langle \mathcal{L}_2^{-1}(1), 1 \rangle) = 1$$

³ By the previous example, we see that such a critical wavespeed also exists for $p = 4$.

and, furthermore, that at the nonlinear center we have

$$\begin{aligned} D_{\text{gKdV}}(a = 0, E^*(c), c, b = 0) &= -\Omega(c, p)V'' \left((1 - c^2)^{1/p}, 0, 1 - c^2 \right)^{-9/2} \\ &= -\Omega(c, p) \left((1 - c^2)p \right)^{-9/2} \end{aligned}$$

for some positive constant $\Omega(c_0, p) > 0$ (see [19, Section 5] for details). It follows that in a neighborhood of the equilibrium solution $u \equiv (1 - c^2)^{1/p}$ there exists a critical wavespeed $c(p, E)$ such that all nearby small-amplitude periodic traveling wave solutions of (4.9) of the form $u(\cdot; a = 0, E, c, b = 0)$ with $E_0 < E \ll 1$ are orbitally stable for $c^*(p, E) < |c| < 1$, and are spectrally unstable to perturbations with $k_r = 1$ for $0 < |c| < c^*(p, E)$. In particular, the equilibrium solution itself is orbitally stable for all $|c| \in (-1, 1)$, implying that $c^*(p, E) \rightarrow 0^+$ as $E \rightarrow (E_0^*)^-$, i.e. as one approaches the equilibrium solution. Note this is consistent with the numerical calculations of $c^*(p, E)$ presented in [16] in the cases $f(u) = u^2$ and $f(u) = u^3$.

Remark 4.10. It is interesting to note that when $p > 4$ the waves with $a = b = 0$ undergo a transition to instability as one moves from a neighborhood of the equilibrium solution $u \equiv (1 - c^2)^{1/p}$ to a neighborhood of the limiting solitary wave.

Acknowledgements. TK gratefully acknowledges the support of the Jack and Lois Kuipers Applied Mathematics Endowment, a Calvin Research Fellowship, and the National Science Foundation under grants DMS-0806636 and DMS-1108783. MJ was partially supported by the University of Kansas General Research Fund allocation 2302278. JB gratefully acknowledges support from the National Science Foundation under grant DMS-0807584.

References

1. Angulo, J., Bona, J., Scialom, M.: Stability of cnoidal waves. *Adv. Differ. Equ.* **11**, 1321–1374 (2006)
2. Angulo, J., Quintero, J.R.: Existence and orbital stability of cnoidal waves for a 1D Boussinesq equation. *Int. J. Math. Math. Sci.* **2007**, Article ID 52020 (2007). doi:[10.1155/2007/52020](https://doi.org/10.1155/2007/52020)
3. Arruda, L.: Nonlinear stability properties of periodic travelling wave solutions of the classical Korteweg–de Vries and Boussinesq equations. *Port. Math.* **66**(2), 225–259 (2009)
4. Bona, J., Sachs, R.: Global existence of smooth solutions and stability of solitary waves for a generalized Boussinesq equation. *Commun. Math. Phys.* **118**, 15–29 (1988)
5. Bronski, J., Johnson, M.: The modulational instability for a generalized KdV equation. *Arch. Ration. Mech. Anal.* **197**(2), 357–400 (2010)
6. Bronski, J., Johnson, M., Kapitula, T.: An index theorem for the stability of periodic traveling waves of KdV type. *Proc. R. Soc. Edinb. Sect. A* **141**(6), 1141–1173 (2011)
7. Chugunova, M., Pelinovsky, D.: On quadratic eigenvalue problems arising in stability of discrete vortices. *Linear Algebra Appl.* **431**, 962–973 (2009)
8. Deconinck, B., Kapitula, T.: On the spectral and orbital stability of spatially periodic stationary solutions of generalized Korteweg–de Vries equations. submitted
9. Deconinck, B., Kapitula, T.: The orbital stability of the cnoidal waves of the Korteweg–de Vries equation. *Phys. Lett. A* **374**, 4018–4022 (2010)
10. Fang, Y., Grillakis, M.: Existence and uniqueness for Boussinesq type equations on a circle. *Commun. Part. Differ. Equ.* **21**(7–8), 1253–1277 (1996)
11. Farah, L., Scialom, M.: On the periodic “good” Boussinesq equation. *Proc. Am. Math. Soc.* **138**(3), 953–964 (2010)
12. Grillakis, M.: Analysis of the linearization around a critical point of an infinite dimensional Hamiltonian system. *Commun. Pure Appl. Math.* **43**, 299–333 (1990)
13. Grillakis, M., Shatah, J., Strauss, W.: Stability theory of solitary waves in the presence of symmetry, I. *J. Funct. Anal.* **74**, 160–197 (1987)
14. Grillakis, M., Shatah, J., Strauss, W.: Stability theory of solitary waves in the presence of symmetry, II. *J. Funct. Anal.* **94**, 308–348 (1990)

15. Gurski, K., Kollár, R., Pego, R.: Slow damping of internal waves in a stably stratified fluid. *Proc. R. Soc. Lond. A* **460**, 977–994 (2004)
16. Hakkaev, S., Stanislavova, M., Stefanov, A.: Linear stability analysis for periodic traveling waves of the Boussinesq equation and the KGZ system. *Proc. Roy. Soc. Edinburgh A* (to appear)
17. Hărăguș, M., Gallay, T.: Orbital stability of periodic waves for the nonlinear Schrödinger equation. *J. Dyn. Differ. Equ.* **19**, 825–865 (2007)
18. Hărăguș, M., Kapitula, T.: On the spectra of periodic waves for infinite-dimensional Hamiltonian systems. *Physica D* **237**(20), 2649–2671 (2008)
19. Johnson, M.: Nonlinear stability of periodic traveling wave solutions of the generalized Korteweg–de Vries equation. *SIAM J. Math. Anal.* **41**(5), 1921–1947 (2009)
20. Johnson, M.: On the stability of periodic solutions of the generalized Benjamin–Bona–Mahony equation. *Physica D* **239**(19), 1892–1908 (2010)
21. Kapitula, T.: On the stability of N -solitons in integrable systems. *Nonlinearity* **20**(4), 879–907 (2007)
22. Kapitula, T.: The Krein signature, Krein eigenvalues, and the Krein Oscillation Theorem. *Indiana Univ. Math. J.* **59**, 1245–1276 (2010)
23. Kapitula, T., Promislow, K.: *Spectral and dynamical stability of nonlinear waves*. Springer, New York (2013)
24. Kapitula, T., Promislow, K.: Stability indices for constrained self-adjoint operators. *Proc. Am. Math. Soc.* **140**(3), 865–880 (2012)
25. Kapitula, T., Kevrekidis, P., Sandstede, B.: Counting eigenvalues via the Krein signature in infinite-dimensional Hamiltonian systems. *Physica D* **195**(3&4), 263–282 (2004)
26. Kapitula, T., Kevrekidis, P., Sandstede, B.: Addendum: counting eigenvalues via the Krein signature in infinite-dimensional Hamiltonian systems. *Physica D* **201**(1&2), 199–201 (2005)
27. Kollár, R.: Homotopy method for nonlinear eigenvalue pencils with applications. *SIAM J. Math. Anal.* **43**(2), 612–633 (2011)
28. Kostenko, A.: On the defect index of quadratic self-adjoint operator pencils. *Math. Notes* **72**(2), 285–290 (2002)
29. Liu, Y.: Instability of solitary waves for generalized Boussinesq equations. *J. Dyn. Differ. Equ.* **5**, 537–558 (1993)
30. Lyong, N.: The spectral properties of a quadratic pencil of operators. *Russ. Math. Surv.* **48**, 180–182 (1993)
31. Markus, A.: Introduction to the spectral theory of polynomial operator pencils. In: *Translation of Mathematical Monographs*, vol. 71. American Mathematical Society, Providence (1988)
32. Pelinovsky, D.: Inertia law for spectral stability of solitary waves in coupled nonlinear Schrödinger equations. *Proc. R. Soc. Lond. A* **461**, 783–812 (2005)
33. Pivovarchik, V.: On spectra of a certain class of quadratic operator pencils with one-dimensional linear part. *Ukr. Math. J.* **59**(5), 766–781 (2007)
34. Shkalikov, A.: Operator pencils arising in elasticity and hydrodynamics: the instability index formula. In: Gohberg, I., Lancaster, P., Shivakumar, P. (eds.) *Recent Developments in Operator Theory and its Applications*. Operator Theory Advances and Applications, vol. 87, pp. 358–385 (1996)
35. Stanislavova, M., Stefnov, A.: Linear stability analysis for traveling waves of second order in time PDE's. *Nonlinearity* **25**(9), 2625–2654 (2012)

Communicated by P. Constantin